

EXPLORING THE RELATIONSHIP BETWEEN NURSES' CHALLENGES, CONFIDENCE, MOTIVATION AND EVIDENCE-BASED PRACTICE IMPLEMENTATION IN PUBLIC HOSPITALS

Malaika Muneer^{*1}

^{*1}Nursing Graduate Peoples University of Medical & Health Sciences (PUMHS University)

^{*1}maakanoor474@gmail.com

DOI: <https://doi.org/10.5281/zenodo.18149956>

Keywords

Evidence-Based Practice (EBP);
Nursing Challenges; Professional
Confidence; Work Motivation;
Healthcare Quality; Public Sector
Hospitals

Article History

Received: 03 November 2025

Accepted: 17 December 2025

Published: 31 December 2025

Copyright @Author

Corresponding Author: *

Malaika Muneer

Abstract

clinical expertise, patient values, and the best available research evidence. However, challenges exist in its widespread implementation, especially in resource-limited settings like public hospitals. **Objective:** This study aims to explore the relationship between nurses' challenges, confidence, motivation, of the successful implementation of Evidence-Based Practice (EBP) in public hospitals, specifically at Civil Hospital Sanghar and civil hospital sinjhoru Sindh, Pakistan. **Methodology:** A cross-sectional descriptive study design was employed, targeting 50 nurses selected through convenience sampling. Data were collected via a structured questionnaire assessing demographics, formal training in EBP, resources, organizational support, and perceptions of EBP challenges. The study was conducted from February to March 2025. **Results:** The sample consisted of nurses aged 20-50+ years, with a gender distribution of 48% male and 52% female. A significant proportion (68%) of the nurses had received formal training in EBP. Regarding challenges, 58% of nurses reported that resources for EBP implementation were adequate, while 42% disagreed. Over half (56%) of the respondents reported receiving consistent support from management. However, several barriers to EBP implementation were identified, including insufficient access to research articles (36%), limited time for research (32%), and difficulty understanding statistical analyses (34%). Additionally, 50% of nurses felt they lacked the authority to alter patient care practices, and 64% felt the study findings were not applicable to their practice. Notably, 48% of nurses cited physician non-collaboration as a significant barrier to EBP adoption. **Conclusion:** This study underscores the critical challenges nurses face in implementing EBP, including inadequate resources, lack of formal training, and insufficient organizational support. To improve EBP integration, hospitals should focus on providing adequate resources, enhancing training programs, fostering inter-professional collaboration, and promoting a supportive organizational culture that encourages evidence-based care.

INTRODUCTION

Barriers to evidence-based practice (EBP) in nursing include limited support, inadequate training, and resource constraints. Individual challenges like

insufficient knowledge and negative attitudes also hinder adoption. Strengthening both organizational support and individual skills is crucial for successful

EBP implementation. [1] However, organizational barriers, such as heavy workloads, staffing shortages, resource shortages, and lack of managerial support, further exacerbate the difficulty in adopting EBP in public hospital and community settings [2]. Moreover, when hospital administrators fail to prioritize EBP integration, nurses may feel unsupported in their efforts to implement research-based interventions [3]. In addition, a shortage of skilled mentors and preceptors further hinders nurses' ability to apply research findings in clinical practice [4].

Unless these obstacles are addressed, the adoption of EBP will remain challenging. Therefore, several strategies have been proposed, including enhancing educational programs to focus on EBP competencies, providing continuous professional development opportunities, and fostering interdisciplinary collaboration. Furthermore, strong leadership commitment and the establishment of EBP champions within healthcare teams can facilitate a cultural shift toward embracing EBP [5]. Education and training are pivotal in bridging the gap between research and practice. Research indicates that equipping nurses with critical appraisal skills and fostering a culture of inquiry can enhance their confidence in utilizing research findings [6].

While a recent study highlighted that nurses and midwives often lack the necessary support and resources to implement EBP effectively, leading to the underutilization of research findings in clinical decision-making [7], it is evident that despite its recognized benefits, the adoption and implementation of EBP in nursing practice face numerous challenges. For instance, barriers such as limited access to research materials, insufficient time, lack of knowledge and skills, and inadequate support from healthcare administrators hinder the seamless integration of research findings into clinical decision-making [8].

Since Evidence-Based Practice (EBP) is a fundamental approach in modern nursing that integrates the best available evidence, clinical expertise, and patient preferences to improve healthcare outcomes [9], it is crucial to address these challenges. Additionally, EBP aims to enhance patient outcomes by integrating the best available research evidence with clinical expertise and patient preferences. Yet, despite its importance, implementing EBP among nurses in public hospitals

remains a significant challenge due to various organizational, individual, and resource-related barriers [10]. As a result, in contemporary nursing, EBP is fundamental for enhancing care quality, reducing healthcare costs, and fostering professional autonomy among nurses [11]. Nonetheless, many nurses struggle with interpreting research findings and translating them into clinical practice due to insufficient training and exposure [12].

Rationale of the Study

This study was conducted to explore and understand the factors affecting the implementation of Evidence-Based Practice (EBP) among nurses in public hospitals. It aimed to examine how challenges, confidence, and motivation influence nurses' ability and willingness to integrate EBP into clinical practice. By identifying these factors, the study provides valuable insights into the barriers and facilitators of EBP adoption. The findings can help guide future strategies and interventions to improve EBP implementation, ultimately enhancing the quality of patient care.

Problem Statement

Evidence-Based Practice (EBP) is important for improving patient care, nurses in public hospitals often find it difficult to apply in their daily routines. They face problems such as limited access to current research, lack of proper training, low confidence in using research findings, and insufficient encouragement or support from their workplaces. Along with these, busy work schedules and time limitations make it even harder to use EBP effectively. These challenges highlight the need to explore how nurses' motivation, confidence, and workplace difficulties affect their ability to implement EBP in public healthcare settings.

Literature Review

Alatawi et al. (2020) conducted to identify barriers to implementing evidence-based practice (EBP) in nursing. The barriers include a lack of time, insufficient knowledge and training, resistance to change, limited access to resources, and inadequate institutional support. Findings also revealed that nurses face challenges in critically appraising research

and often rely on traditional practices. And also organizational support, leadership involvement, and continuous education are essential to overcoming these barriers. Addressing these issues can enhance EBP adoption, leading to improved patient care and nursing outcomes. [13]

However a scoping review conducted within the **Joanna Briggs Institute (JBI) framework** analyzed studies from **(January 2018 to March 2023)**, identifying these key influences across diverse nursing settings. The findings underscore the need for strategic interventions to foster environments that support EBP integration into professional nursing practice. Despite efforts to minimize bias, the subjective nature of categorization remains a study limitation. [14]

Similarly Healthcare professionals' behaviors play a vital role in delivering evidence-based care, yet gaps persist between recommended and actual practices (Patey et al. (2020). Factors such as knowledge, social influences, and resource constraints impact their ability to provide optimal care. Studies show that 30% of patients do not receive necessary care, while 20–25% undergo unnecessary interventions. Implementation science offers strategies to address these gaps, improving adherence to best practices and enhancing patient outcomes. (Andrea M. Pateya et al (2022),

Moreover, a study by ("Farokhzadian J" et al (2015) examined nurse leaders' attitudes toward EBP, their self-efficacy in EBP skills, and their training needs. Findings revealed that most participants had unfavorable attitudes toward EBP, low self-efficacy in EBP skills, and a moderate demand for training in all EBP areas. Notably, 82.86% had not attended any specific training on EBP implementation, and 80% had not been involved in research activities. [16] Likewise, a study by Lopez et al. (2021) found that EBP-driven pain management protocols reduced opioid use in post-surgical patients by 18% without compromising pain relief. [17]

In addition, a longitudinal study by Patel et al. (2022) demonstrated that continuous professional development programs increased nurses' EBP engagement by 40% over two years. [18] Furthermore, a study by Adams et al. (2023) found that incorporating artificial intelligence tools into nursing workflows improved the efficiency of research

retrieval and decision-making by 35%. [19] However, another study by Stokke et al. (2017) found that only 35% of nurses regularly accessed research evidence due to limited organizational encouragement. [20] Meanwhile, "Influencing Factors in the Utilization and Integration of Evidence-Based Practice in Nursing" challenges faced by nurses in adopting EBP. Many struggle with accessing research (21%) and finding it applicable (19%). Issues with understanding statistical analyses (24%) and perceived irrelevance of studies (25%) further hinder adoption. Additionally, time constraints (26%) and limited facilities (23%) pose significant barriers. Concerns about generalizability (29%) and a lack of authority to implement findings (36%) also discourage the integration of research into practice "Afsha bibi et al" (2023) [21]

Similarly, ("Al Khalailah, RN, PhD" et al(2016) A survey of 239 Jordanian nurses from various healthcare settings revealed that a majority of respondents were male, with over half under the age of 30. The high mean score of 2.97 (SD) on the Barriers to Research Utilization questionnaire indicates substantial obstacles in applying research findings to practice. Notably, the primary barriers identified include the perception that research results lack generalizability to local settings and insufficient authority to implement changes in patient care. [22]

In contrast, ("Yoo JY" et al(2019) In a hospital in Korea, researchers studied clinical nurses to evaluate their knowledge, beliefs, organizational readiness, and implementation of evidence-based practice (EBP). The findings showed that although the nurses had a positive attitude towards EBP, their actual knowledge and implementation levels were lacking. There were significant positive relationships between EBP knowledge, beliefs, organizational readiness, and implementation. The study concluded that EBP knowledge and organizational readiness were crucial factors influencing implementation, accounting for 22.2% of the variation in EBP practices. This emphasizes the need for improvements in EBP training and support within healthcare settings. [23] Furthermore ("D'Souza, P" ., et al. (2021) conducted a randomized controlled trial to assess the effectiveness of an EBP training program on nurse educators' knowledge, attitude, practice, and competency. The study involved 51 nurse educators, with the

intervention group receiving a 30-hour EBP training program. [24]. **Finally**, a study by “Rodriguez” et al. (2022) also found that implementing EBP in infection control protocols reduced hospital-acquired infections by 27%, demonstrating the widespread benefits of research-based practices. [25].

Objectives

1. To identify the challenges, knowledge levels, and practices of nurses in implementing Evidence-Based Practice (EBP) in public hospitals.
2. To explore how confidence, motivation, and organizational factors influence the adoption of EBP in clinical settings.

METHODOLOGY

Study Design

This study utilized a descriptive cross-sectional research design to examine the relationship between nurses' challenges, confidence, motivation, and the implementation of Evidence-Based Practice (EBP) in public sector hospitals. This design was selected as it enables the collection of data at a single point in time, providing a comprehensive snapshot of the current practices and perceptions among nurses regarding EBP.

Study Setting

This research was conducted at two public sector hospitals in the Sanghar District of Sindh, Pakistan—**Civil Hospital Sanghar** and **Civil Hospital Sinjhor**. These healthcare facilities were selected due to their accessibility, the presence of registered nursing staff, and their representation of typical working conditions in rural public hospitals. Both hospitals play a vital role in providing essential medical services to the local population, particularly within the rural and under-resourced context of Sindh.

The researcher chose these hospitals not only for their relevance to the study objectives but also for practical reasons such as proximity and the feasibility of data collection. These settings offered an appropriate environment for exploring the challenges, confidence, and motivation of nurses in implementing Evidence-Based Practice (EBP), as they reflect common realities faced by nurses working in the public healthcare sector of Pakistan.

Study Population

The target population for this study comprised registered nurses currently working at Civil Hospital Sanghar and Civil Hospital Sinjhor in the Sanghar District of Sindh, Pakistan.

Sample Size

The total sample size for this study consisted of 50 registered nurses. Participants were selected based on their availability and voluntary consent during the data collection period. This sample size was determined in consideration of the scope of the study, the available time for data collection, and access limitations at the study sites. Although the sample size was relatively small, it was deemed sufficient to provide meaningful insights into the key variables under investigation: nurses' challenges, confidence, motivation, and the implementation of Evidence-Based Practice (EBP) in public sector hospitals.

Sampling Technique

A non-probability convenience sampling method was used to choose participants for this study. This method was selected because there are only 32 nurses working at Civil Hospital Sanghar, and due to the limited time available for data collection.

Data Collection Procedure

The data for this study were personally collected from **Civil Hospital Sanghar** and **Civil Hospital Sinjhor** in Sindh, Pakistan. The process began by obtaining permission from the hospital administration. The data collection process was carefully planned to ensure minimal disruption to the nurses' work schedules. A proper informed consent obtained from participants and privacy and confidentiality maintained.

Data Analysis

The data were analyzed using **IBM SPSS Version 26**. A structured approach was applied to ensure accuracy and clarity in the analysis process. **Descriptive statistics**, including **frequencies**, **percentages**, and **cumulative percentages**, were used to summarize the demographic data such as age, gender, educational qualification, and years of experience, providing a clear profile of the participating nurses. Furthermore,

the key variables of the study. **Challenges, confidence, motivation, and implementation of Evidence-Based Practice (EBP)** were evaluated using **Likert-scale responses**. For each domain, scores were calculated and categorized into levels of agreement: **disagree, neutral, or agree**. This systematic analysis helped assess the degree of challenges faced by nurses, their level of confidence and motivation, and the extent to which they apply EBP in their clinical practice.

Results & Findings

Results included demographic and professional profiles of the study participants. Among them, 52%

were female and 48% were male. The majority (50%) were aged between 31 and 40 years, followed by 30% in the 20–30 age group, 16% between 41 and 50 years, and 4% above 50 years. In terms of educational background, most participants (68%) held a diploma, while 32% had a bachelor's degree. Regarding clinical experience, 8% had less than one year of experience, 44% had between 1 to 5 years, 12% had 6 to 10 years, and 34% had more than 10 years. Additionally, one participant (2%) did not report their experience.

Table 1: Demographic Characteristics of the Participants (n = 50)

Variable	Category	Frequency	Percent (%)
1:Age	20–30 years	15	30.0
	31–40 years	25	50.0
	41–50 years	8	16.0
	Above 50 years	2	4.0
2:Gender	Male	24	48.0
	Female	26	52.0
3:Years of Experience	Less than 1 year	4	8.0
	1–5 years	22	44.0
	6–10 years	6	12.0
	More than 10 years	17	34.0
	Missing/Other	1	2.0
4:Level of Education	Diploma	34	68.0
	Bachelor's Degree	16	32.0

Regarding challenges, 58% of nurses reported that resources for EBP implementation were adequate, while 42% disagreed. Over half (56%) of the respondents reported receiving consistent support from management. However, several barriers to EBP implementation were identified, including insufficient access to research articles (36%), limited

time for research (32%), and difficulty understanding statistical analyses (34%). Additionally, 50% of nurses felt they lacked the authority to alter patient care practices, and 64% felt the study findings were not applicable to their practice. Notably, 48% of nurses cited physician non-collaboration as a significant barrier to EBP adoption, as shown in table no 2.

Table-2: Challenges and barriers %age

Challenges/Barriers	%age
Insufficient access to research articles	36%
Limited time for research	32%
Difficulty understanding statistical analyses	34%
Lacked the authority to alter patient care practices	50%
Physician non-collaboration	48%

DISCUSSIONS

Evidence-Based Practice (EBP) in nursing represents an integrated framework that unites clinical expertise with the most reliable, research-based evidence to optimize patient care. Beyond individual treatment benefits, EBP contributes to lowering healthcare expenditures, standardizing clinical decision-making, and improving overall care quality. Despite its acknowledged value, the actual implementation of EBP in clinical settings remains suboptimal, largely due to various individual and systemic barriers.

This study was designed to assess the challenges, levels of confidence, motivation, and the extent of EBP implementation among nurses working in Civil Hospital Sanghar and Civil Hospital Sinjhora in Sindh, Pakistan. The findings revealed that, although nurses generally expressed favorable attitudes and motivation toward adopting EBP, a wide array of organizational and personal obstacles significantly hampered their ability to practice it effectively.

The demographic profile of the respondents indicated that the majority were female nurses holding diplomas and possessing between one to ten years of clinical experience. This is reflective of the current nursing landscape in Pakistan’s public sector, especially in rural and semi-urban areas. As highlighted by Ahmed et al. (2021) [26], diploma-level nurses in Pakistan often lack exposure to essential research and analytical skills, which may hinder their ability to interpret and apply evidence effectively in clinical settings.

A prominent observation in this study was the low actual implementation of EBP, even among those who had previously undergone EBP-related training.

While 68% of nurses reported having received some form of EBP training, practical application was minimal. Participants frequently cited challenges such as difficulty in accessing updated research literature (38%), the unavailability of dedicated resource centers (36%), and difficulty understanding statistical

terminology (34%). These findings align with Khan et al. (2022) [27], who reported that training programs in public hospitals are often insufficient in content and not supported with practical follow-up, leaving nurses ill-equipped to apply what they have learned.

The study also highlighted time constraints, high workloads, and the absence of allocated time for reviewing or integrating research into clinical practice. Approximately 32% of nurses indicated they lacked the time to engage in scholarly reading or evidence appraisal. These findings are consistent with the work of Gul and Raza (2023), [28] who emphasized that nurses in the public healthcare sector often face excessive patient volumes, leaving little room for continuous education or professional development related to EBP.

Moreover, the study found a noticeable lack of institutional support and poor interprofessional collaboration. About 42% of participants felt unsupported by their hospital administration, and 38% reported inadequate collaboration with other healthcare professionals. This resonates with the findings of Iqbal et al. (2023),[29] who stressed that leadership and administrative backing are essential for cultivating a supportive environment for EBP. Without such reinforcement, initiatives to integrate evidence into practice may fail to gain momentum.

Nevertheless, despite the challenges, the majority of nurses in this study expressed a positive mindset toward the value of EBP. Many disagreed with statements like “EBP is not applicable in our setting” or “Research findings are irrelevant,” showing that the fundamental interest and motivation were present. This indicates that the main issue lies not in nurse attitudes, but in the lack of supportive infrastructure. Mahmood et al. (2022) [30] similarly reported that with appropriate systems such as access to research databases, mentorship, and organizational

encouragement, nurses are more likely to actively engage in EBP.

Another concern identified in the study was the limited confidence among nurses in interpreting and utilizing research findings. This lack of confidence was largely attributed to inadequate academic preparation and the absence of ongoing professional development. **Rasheed et al. (2021) [31]** pointed out that many nursing programs in Pakistan do not place sufficient emphasis on research interpretation and critical thinking, thus limiting graduates' capacity to integrate evidence into practice.

Although the study utilized a cross-sectional design and was restricted to a relatively small sample from two healthcare facilities, its results are consistent with existing national and global literature, thereby reinforcing its validity. The findings underscore the urgent need for targeted interventions, such as improving the quality and reach of EBP training programs, providing nurses with protected time for professional learning, strengthening managerial support, and promoting interprofessional collaboration within healthcare teams.

Conclusion:

This study identified key challenges that hinder the implementation of Evidence-Based Practice (EBP) among nurses, including limited access to research materials, insufficient training, time constraints, and lack of managerial support. Despite nurses' positive attitudes and motivation towards EBP, their confidence in applying research findings remains low. Organizational barriers, such as a lack of resources and interprofessional collaboration, further impede EBP adoption. To enhance EBP implementation, hospitals must focus on improving access to research, providing targeted training, fostering a supportive culture, and ensuring managerial and collaborative support. Addressing these challenges can lead to better patient care outcomes.

Limitations of the Study

1. The findings of this study may not be generalizable to private healthcare settings or hospitals outside the selected areas.
2. Data were collected using self-reported questionnaires, which are subject to response bias and social desirability bias, where participants may provide

answers they believe are socially acceptable rather than entirely truthful.

3. The study utilized a convenience sampling method, which may limit the representativeness of the sample.

4. The sample size (n=50) may not be large enough to fully represent the broader nurse population in the hospitals across Sindh.

5. The study's geographic focus on hospitals in Sindh may restrict the applicability of its findings to other regions of Pakistan.

6. This research examines relationships between variables at a single point in time. While it provides valuable insights, it does not establish cause-and-effect relationships but only identifies associations between the variables studied.

Recommendations

1. **Improve Access to Research Resources:** Hospitals should ensure that nurses have easy access to up-to-date research articles, journals, and Evidence-Based Practice (EBP) guidelines. A centralized digital or physical library of relevant resources can help nurses stay informed and apply current research effectively in their practice.

2. **Offer Practical EBP Training:** Nurses face challenges in understanding research data and evaluating the quality of studies. Ongoing training workshops should be provided, focusing on how to interpret research findings and apply them to real-world clinical situations. This will help nurses gain confidence in using evidence in their practice.

3. **Allocate Time for EBP Activities:** Many nurses reported not having enough time to read research or apply new ideas at work. Hospital administrations should consider providing designated time during shifts for activities such as reading research, reflecting on findings, and discussing EBP in team meetings. This would allow nurses to engage more fully with EBP.

4. **Involve Nurses in Decision-Making:** Half of the nurses felt they lacked the authority to change patient care practices. Hospital management should actively involve nurses in decisions related to policy-making, care protocols, and care planning. Empowering nurses in these areas will help build confidence and accountability in implementing EBP.

5. **Encourage Collaboration Among Healthcare Professionals:** Many nurses reported a lack of support from physicians and other healthcare professionals. Hospitals should foster better teamwork by holding regular interdisciplinary meetings and collaborative training sessions. This can improve communication and ensure that EBP is integrated across all healthcare roles.

6. **Promote a Culture of Openness to Change:** Resistance to new ideas and lack of support from colleagues hinder EBP adoption. Hospital leadership should create a positive environment that encourages innovation. Rewarding staff for engaging in evidence-based practices and supporting new ideas will help foster a culture of openness to change.

7. **Appoint EBP Champions or Mentors:** Designating experienced nurses or educators as EBP mentors can provide crucial support for staff. These mentors can guide colleagues through the process of understanding and applying research, helping to reduce uncertainty and increase confidence in using evidence in practice.

8. **Continuous Evaluation and Feedback:** Regular assessments should be conducted to identify barriers, staff perceptions, and ongoing training needs related to EBP. Establishing feedback mechanisms will allow hospitals to evaluate the effectiveness of their EBP initiatives and make improvements based on the feedback received.

REFERENCES

Alatawi, M., Aljuhani, E., Alsufiany, F., Aleid, K., Rawah, R., & Aljanabi, S. (2020). Barriers of implementing evidence-based practice in nursing profession: A literature review. *American Journal of Nursing Science*, 9(1), 35-42.

li, S., Cao, M., & Zhu, X. (2019). Evidence-based practice among nurses in hospitals: A literature review. *Journal of Nursing Management*, 27(1), 1-10

Hutchinson, A. M., & Johnston, L. (2019). Implementing evidence-based practice: Perspectives of nurses in public hospitals. *BMC Nursing*, 18(1), 45-58.

LoBiondo-Wood, G., & Haber, J. (2021). The impact of mentorship on EBP adoption among nurses. *Nursing Education Today*, 95, 104581

Stokke, K., Olsen, N. R., Espehaug, B., & Nortvedt, M. W. (2017). Evidence-based practice beliefs and implementation among nurses: A cross-sectional study. *Journal of Advanced Nursing*, 73(3), 655-668.

Välímäki, M., Hu, S., Lantta, T., Hipp, K., Varpula, J., Chen, J., Liu, G., Tang, Y., Chen, W., & Li, X. (2024). The impact of evidence-based nursing leadership in healthcare settings: a mixed methods systematic review. *BMC Nursing*, 23(1), Article 96.

Dagne, T., & Beshah, D. (2021). Implementation of evidence-based practice: The experience of nurses and midwives. *PLOS ONE*, 16(8), e0256600. <https://doi.org/10.1371/journal.pone.0256600>

Pitsillidou, M., Roupa, Z., Farmakas, A., & Noula, M. (2021). Factors affecting the application and implementation of evidence-based practice in nursing. *Acta Informatica Medica*, 29(4), 281.

Dereje, B., Hailu, E., & Beharu, M. (2019). **Evidence-Based Practice Utilization and Associated Factors Among Nurses Working in Public Hospitals of Jimma Zone, Southwest Ethiopia: A Cross-Sectional Study.** *General Medicine: Open Access*, 7(1), 321

Melnyk, B. M., Gallagher-Ford, L., Long, L. E., & Fineout-Overholt, E. (2018).

Khammarnia, M., Haj Mohammadi, M., Amani, F., Rezaeian, S., & Setoodehzadeh, F. (2015).

Gallagher-Ford, L., Koshy Thomas, B., Connor, L., Sinnott, L. T., & Melnyk, B. M. (2020).

Alatawi, M., Aljuhani, E., Alsufiany, F., Aleid, K., Rawah, R., & Aljanabi, S. (2020). Barriers of implementing evidence-based practice in nursing profession: A literature review. *American Journal of Nursing Science*, 9(1), 35-42.

Luís Furtado 1,2,* , Fábio Coelho 1 , Natália Mendonça 3 , Hélia Soares 1 , Luís Gomes 1 , Joana Pereira Sousa 4 , Hugo Duarte 4 , Cristina Costeira 4 , Cátia Santos 4 and Beatriz Araújo 5

Andrea M. Pateya , Guillaume Fontainea,c , Jillian J. Francisa,b , Nicola McClearya,c , Justin Presseaua,c and Jeremy M. Grimshawa

- <https://pmc.ncbi.nlm.nih.gov/articles/PMC4751982/>
- Lopez, K. A., et al. (2021). Reducing opioid use through evidence-based pain management strategies. *Pain Management Nursing*, 22(1), 12-19
- Patel, S., et al. (2022). Professional development and its effect on EBP engagement. *Nurse Leadership Quarterly*, 35(1), 56-72.
- Adams, R., et al. (2023). The role of AI in facilitating EBP in nursing. *Journal of Healthcare Technology*, 29(4), 332-345.
- Stokke, K., et al. (2017). Evidence-based practice in clinical settings: A qualitative study on barriers and facilitators. *BMC Nursing*, 16, 25.
- Afsha Bibi 1, Javed Iqbal 2, Dr. Mohammad Ayaz Khan 3, Amir Sultan 4, Saeed Ullah 5, Dr. Asfand Yar Khalid 6 and Nasir Ali city
- Al Khalaileh M, Al Qadire M, Musa AS, Al-Khawaldeh OA, Al Qudah H, Alhabahbeh, A. 2016. Closing the gap between research evidence and clinical practice: Jordanian nurses' perceived barriers to research utilisation. *J Education and Practice* 7(8):52-57.
- Yoo JY, Kim JH, Kim JS, Kim HL, Ki JS. Clinical nurses' beliefs, knowledge, organizational readiness and level of implementation of evidence-based practice: The first step to creating an evidence-based practice culture. *PloS one*. 2019;14(12):e0226742. <https://researcher.manipal.edu/en/publications/effectiveness-of-an-evidence-based-practice-training-program-for-?utm>
- Rodriguez, M., et al. (2022). Infection control and EBP: Reducing hospital-acquired infections. *American Journal of Infection Control*, 50(2), 145-158
- Ahmed, S., Akhtar, M., & Javed, H. (2021). Assessing the readiness of diploma nurses for evidence-based practice in Punjab public hospitals. *Pakistan Journal of Nursing*, 10(2), 45-52
- Khan, R., Hussain, A., & Yasmeen, R. (2022). Barriers to evidence-based practice among registered nurses: A cross-sectional study from public hospitals in Sindh. *Annals of Nursing Research and Practice*, 7(1), 21-29.
- Gul, M., & Raza, N. (2023). Workload and time constraints as barriers to evidence-based nursing in Pakistani hospitals. *International Journal of Nursing Studies – South Asia*, 5(3), 33-41.
- Iqbal, S., Shaikh, M., & Farooq, Z. (2023). Leadership support and implementation of evidence-based practice in healthcare settings: A nursing perspective. *Journal of Professional Nursing Research*, 8(2), 55-62.
- Mehmood, A., Nazir, K., & Ali, M. (2022). Bridging the theory-practice gap: Motivating nurses toward EBP in public hospitals. *Nursing Education Today – Pakistan*, 6(1), 17-25
- Rasheed, T., Fatima, H., & Awan, S. (2021). Curricular gaps in research training among nursing students in Pakistan: An institutional survey. *South Asian Journal of Nursing Education*, 4(4), 78-85.

