The Research of Medical Science Review
ISSN: 3007-1208 & 3007-1216 Volume 4, Issue 1, 2026

DENGUE FEVER IN PAKISTAN: A COMPREHENSIVE REVIEW OF EPIDEMIOLOGY, CLINICAL PRESENTATION, PATHOPHYSIOLOGY, AND MANAGEMENT

Saba Malik[1], Hafiza Sadia[2], Ahmad Ali[3], Sheeba Riaz[4], Muhammad Naeem[5], Iqra Bibi[6], Ghulam Shabbir[7], Muhammad Sikandar Majid[*8]

[1,3, 6,7,*8] *Department of Zoology, Wildlife and Fisheries, University of Agriculture, Faisalabad, Pakistan*
[2]*Department of Plant Breeding and Genetics, University of Agriculture, Faisalabad, Pakistan*
[4]*School of Biochemistry and Biotechnology, University of the Punjab, Lahore, Pakistan*
[5]*College of Biological Resources and Food Engineering, Qujing Normal University, Qujing, China.*

[1]majidsikandar2@gmail.com

[1]https://orcid.org/0000-0002-8796-1706, [2]https://orcid.org/0009-0001-4448-2510,
[3]https://orcid.org/0009-0007-2546-9736, [4]https://orcid.org/0009-0000-5763-503X,

DOI: https://doi.org/10.5281/zenodo.18239516

Keywords
Dengue fever, Pakistan, Epidemiology, Clinical presentation, Pathophysiology, Aedes mosquito, Dengue hemorrhagic fever, Vector control, Disease management, Public health, Prevention strategies, Outbreak response

Article History
Received: 20 November 2025
Accepted: 28 December 2025
Published: 13 January 2026

Copyright @Author
Corresponding Author: *
Muhammad Sikandar Majid

Abstract

Dengue fever is a mosquito-borne viral disease that poses a significant public health challenge in Pakistan and across South Asia. Transmitted primarily by the female Aedes mosquito, dengue presents with a spectrum of clinical manifestations, ranging from mild febrile illness to severe, life-threatening complications such as Dengue Hemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS). This comprehensive review synthesizes data from published studies conducted in Pakistan between 1988 and 2018, highlighting epidemiological trends, clinical presentations, pathophysiology, and management strategies. Analysis indicates that males (62.4%) are more frequently affected than females (37.6%), with the highest incidence observed in the 21–30-year age group. Common symptoms include high-grade fever, body aches, nausea, and headache, with severe cases showing hemorrhagic manifestations. Effective management relies on a combination of environmental, biological, and chemical interventions, alongside preventive measures such as vaccination, personal protection, and community awareness. Despite ongoing initiatives like the Punjab Health Line Project, gaps in healthcare infrastructure, physician knowledge, and rapid outbreak response remain challenges. This review underscores the urgent need for coordinated strategies encompassing vector control, clinical preparedness, and public education to mitigate the growing burden of dengue in Pakistan.

INTRODUCTION

The term *dengue* is derived from the Spanish word for "fastidious" and is thought to have originated from the Swahili word *dinga*, meaning "evil spirit," reflecting early perceptions of the disease's debilitating nature (Mehmood et al., 2021). The first clinical description of dengue,

referred to as "breakbone fever" due to its severe musculoskeletal pain, was documented by Benjamin Rush in 1789 during an epidemic. According to the World Health Organization (WHO), dengue is a mosquito-borne viral infection with a clinical spectrum ranging from mild febrile illness to severe, life-threatening hemorrhagic forms (Mahajan, 2018). WHO estimates indicate that between 50 and 100 million dengue infections occur annually across more than 100 endemic countries, placing nearly half of the global population at risk. Recent surveillance data from the WHO global dengue monitoring network which includes 103 participating countries reported over 8 million cases and more than 3,000 fatalities between January and April 2024, with the highest incidence recorded in March. However, underreporting remains a major challenge, as 28 countries have yet to provide data, suggesting that the true global burden may be considerably higher (Venkatesan, 2024).




Figure 1. A: Geographical distribution and case/death notifications of dengue reported to WHO from January to April 2024. (Note: The surveillance system is still under development and not all countries are currently reflected.) B. Reported cases and deaths worldwide

(https://www.who.int/emergencies/disease-outbreak-news/item/2024-DON518).

The dengue virus (DENV) is a single-stranded ribonucleic acid (RNA) virus belonging to the Flaviviridae family (Solomon, 2000). Unlike many other RNA viruses, DENV exists in four distinct serotypes DENV-1, DENV-2, DENV-3, and DENV-4 each capable of causing dengue fever or its more severe manifestation, dengue hemorrhagic fever (DHF). Infection with one serotype confers lifelong immunity against that specific type but provides only temporary and partial cross-immunity to the others, which increases the risk of severe disease upon secondary infection (Müller et al., 2017). In addition to systemic symptoms, dengue infection may involve multiple organ systems, including the liver, kidneys, brain, and heart, contributing to clinical complications (Roy et al., 2021). At present, no specific antiviral therapy exists for dengue; management primarily relies on timely fluid and electrolyte replacement to prevent dehydration and shock (Hasan et al., 2016). Major epidemiological shifts in dengue virus transmission patterns have been noted since World War II, driven by urbanization, globalization, and increased vector adaptation (Murugesan & Manoharan, 2020).

The primary aim of this study is to provide a comprehensive overview of dengue fever in Pakistan, encompassing its epidemiology, clinical presentation, pathophysiology, and management strategies. Specifically, the study seeks to analyze historical and recent data on dengue incidence, demographic patterns, and clinical outcomes in Pakistan, while evaluating the effectiveness of current preventive, environmental, biological, chemical, and healthcare-based interventions. By synthesizing national and international research findings, this study aims to identify gaps in knowledge, diagnostic practices, and vector control measures, and to propose evidence-based recommendations to strengthen dengue surveillance, management, and public health response in Pakistan.

1. Etiology of the Dengue fever

The dengue virus (DENV) is an enveloped, 50 nm Flavivirus characterized by icosahedral symmetry and surface envelope (E) glycoproteins that facilitate host cell attachment and membrane fusion (Li & Kang, 2022). Its single-stranded RNA genome encodes both structural (C, M, E) and non-structural (NS1–NS5) proteins essential for viral replication, maturation, and evasion of host immune defenses (Sun & Varani, 2022; Songprakhon et al., 2020). Transmission primarily occurs through the bite of infected female Aedes mosquitoes, particularly Aedes aegypti. When a mosquito feeds on the blood of an infected human typically within 2 to 10 days of symptom onset the virus replicates in the mosquito's midgut cells and subsequently disseminates to other tissues, including the salivary glands, enabling further transmission during subsequent bites (Pourzangiabadi et al., 2024). Globally, dengue incidence continues to rise due to vector expansion, geographical spread, and increased epidemic frequency, with Aedes aegypti serving as the dominant vector and other Aedes species acting as secondary transmitters across tropical and subtropical regions (Dehghani & Kassiri, 2021).

The dengue virus (DENV) primarily targets human immune cells, particularly monocytes and dendritic cells, while in mosquitoes, viral replication begins in the midgut epithelium before disseminating to other tissues, including the salivary glands (Lin et al., 2023). DENV employs a range of cell surface receptors to facilitate entry, such as Hsp70 in mosquitoes and heparan sulfate, Hsp90, and C-type lectin receptors in human cells. The Aedes mosquito, which serves as the primary vector, undergoes both aquatic (larval and pupal) and terrestrial (adult) life stages, with transmission potential developing approximately 7–14 days after ingestion of infected blood. Since direct human-to-human transmission does not occur, vector control remains the cornerstone of dengue prevention (Begum et al., 2019). Globally, Aedes aegypti is recognized as the dominant transmitter,

while Aedes albopictus, along with region-specific species such as A. polynesiensis in the Pacific and A. niveus in the Philippines, also contribute to disease spread (Bhugra, 2020).

Aedes mosquitoes, particularly Aedes aegypti and Aedes albopictus, thrive in close proximity to human dwellings, breeding in stagnant water sources such as containers, discarded waste, and household water storage areas. They commonly rest indoors, including under beds and in bathrooms, and are most active during daylight hours, with peak biting periods occurring at dawn and dusk (Krishna et al., 2024). Dengue virus transmission to humans primarily occurs through the bite of an infected female mosquito, which, after an 8–10-day incubation period, becomes capable of transmitting the virus for the remainder of its life without exhibiting any external signs of infection (Ferreira-de-Lima et al., 2018). Although rare, non-vector routes such as vertical transmission (mother-to-child) and blood transfusion have been documented, with some evidence suggesting transovarial transmission the passage of the virus to offspring via eggs (Berube, 2023).

At the cellular level, DENV enters monocytes, macrophages, and dendritic cells through receptor-mediated endocytosis, employing surface molecules like heparan sulfate and C-type lectins for attachment. Following internalization via clathrin-coated vesicles, the acidic environment of the endosome induces fusion between viral and host membranes, releasing viral RNA into the cytoplasm. The genome is then replicated and translated into a polyprotein within the endoplasmic reticulum (ER), where new virions assemble, mature through the ER-Golgi pathway, and are eventually released via exocytosis (Karisa, 2018).

2. Symptoms Created by Dengue Virus

Dengue fever is a systemic, flu-like viral illness that affects individuals of all ages, though the clinical presentation varies with age. Infants and young children typically exhibit a non-specific febrile illness accompanied by rash, whereas older children and adults often develop the classical dengue syndrome, characterized by high-grade fever, severe headache, retro-orbital pain, myalgia, arthralgia, vomiting, abdominal discomfort, and skin rash (Hasan et al., 2016). Additional manifestations may include petechiae, epistaxis, gastrointestinal bleeding, photophobia, nausea, and chills, with fever generally persisting for five to six days, often followed by a maculopapular rash (Htun et al., 2021). In some patients particularly children the disease may appear as an undifferentiated febrile illness without distinctive features. Severe progression can result in Dengue Hemorrhagic Fever (DHF) or Dengue Shock Syndrome (DSS), both characterized by vascular permeability, plasma leakage, and platelet dysfunction, which may culminate in circulatory failure and life-threatening hemorrhage (Dowd et al., 2023).

Table 1: Clinical Stages of Dengue and Associated Symptoms

Classification	Key Features	Common Symptoms	References
Primary Dengue Infection (Febrile Phase)	Begins after 5–7-day incubation	• Sudden high fever (39–40°C) • Headache, myalgia, arthralgia • Nausea, vomiting, loss of appetite • Mild erythema, conjunctival injection • Facial flushing, rash (maculopapular)	Kok et al., (2023)

		• Mild lymphadenopathy	
Classic Dengue Fever (Early Phase)	Non-specific febrile illness which Lasts 5–7 days	• High fever (can reach 40–41°C) • Myalgia, retro-orbital pain • Changes in taste, diarrhea/constipation • Rash, sore throat, cough (infrequent) • Hepatomegaly (soft/tender), rare jaundice • Febrile seizures in children	Khanam et al., (2022)
Secondary Infection / Critical Phase	Occurs in immune individual	• Fluctuating high fever • Hemorrhagic signs • Plasma leakage • Respiratory distress (in DHF) • Higher risk in elderly, neonates • Diabetics • Obese • Renal failure	Dowd et al., (2023)
Dengue Hemorrhagic Fever (DHF)	Most severe secondary manifestation, Occurs ~7–10 days post-onset	• Fever • Spontaneous bleeding (petechiae, purpura, mucosal bleeding) • Plasma leakage (↑ hematocrit) • Thrombocytopenia • Positive tourniquet test • Abdominal pain • vomiting	Bhatt et al., (2021); Prapty et al., (2023)
Dengue Shock Syndrome (DSS)	Most critical stage of dengue	• Circulatory failure • Cold • Clammy extremities • Altered consciousness • Narrow pulse pressure or hypotension • Weak, rapid pulse	Bhatt et al., (2021); Chauhan et al., (2024)

Table 2: Laboratory Diagnosis of Dengue Virus Infection

Category	Test/Method	Target/Indicator	Key Features	References
Clinical Indicators	Routine hematology	Thrombocytopenia (<100,000/mm³), Leukopenia	Early markers; suggest plasma leakage in DHF/DSS.	WHO (2023a)

		(<4,000/mm³), Hemoconcentration (↑Hct ≥20%)		
Virological Tests	NS1 Antigen Test	NS1 protein (Days 1–5)	High specificity; detects primary/secondary infections.	Butt (2008), WHO (2023b)
	RT-PCR	Viral RNA (Days 1–5)	Identifies serotypes (DENV-1–4); high sensitivity.	Kok et al. (2023)
Serological Tests	IgM ELISA	IgM antibodies (Day 5+)	Indicates recent infection.	WHO (2023b)
	IgG ELISA	IgG antibodies (Day 2+ in secondary)	Fourfold rise in titers confirms infection.	Butt (2008)
Emerging Technologies	CRISPR (SHERLOCK/DETECTR)	Viral RNA	Rapid (30–60 min), high accuracy; field-deployable.	Li et al. (2023)
	Multiplex RDTs (NS1 + IgM/IgG)	NS1 + antibodies	Quick (15–30 min) but lower sensitivity than ELISA/PCR.	Khanam et al. (2022)
Imaging	Ultrasound/X-ray	Pleural effusion, ascites	Detects plasma leakage in severe cases (DHF/DSS).	WHO (2023a)

4. Pathophysiology of Dengue

The pathogenesis of dengue fever involves a complex series of immunopathological events initiated by infection with any of the four dengue virus serotypes (DENV-1 to DENV-4). After transmission through the bite of an infected Aedes mosquito, the virus primarily targets dendritic cells and macrophages, which serve as initial replication sites before viral dissemination through the lymphatic and circulatory systems. A pivotal mechanism underlying severe disease progression is antibody-dependent enhancement (ADE), wherein non-neutralizing antibodies from a prior dengue infection facilitate viral entry into Fc receptor-bearing cells during subsequent infection with a different serotype. This process leads to increased viral replication and the excessive production of pro-inflammatory cytokines, culminating in a cytokine storm that contributes to vascular leakage, plasma extravasation, and severe clinical manifestations characteristic of Dengue Hemorrhagic Fever (DHF) and Dengue Shock Syndrome (DSS).

The non-structural protein 1 (NS1) is a key mediator in dengue pathogenesis, driving endothelial dysfunction and activating the complement system, which in turn contributes to the plasma leakage characteristic of severe dengue. This increased vascular permeability can result in hemoconcentration, thrombocytopenia, and, in extreme cases, life-threatening conditions such as dengue hemorrhagic fever or dengue shock syndrome. Host-related factors play a critical role in determining disease severity, with genetic polymorphisms influencing immune responses and younger individuals in endemic regions being particularly susceptible. The interaction between viral determinants and host immune mechanisms produces a wide spectrum of clinical outcomes, ranging from mild febrile illness to severe hemorrhagic manifestations. Secondary infections are often more severe, primarily due to antibody-dependent enhancement (ADE) and immune dysregulation. Accurate diagnosis remains a major challenge in endemic areas, where multiple febrile illnesses present with overlapping clinical features, highlighting the urgent need for precise and reliable diagnostic tools to differentiate dengue from other tropical infections.

5. Distribution of Dengue in Pakistan

Dengue in Pakistan shows varying severity and age distribution. **Jahan (2011)** reported 881 confirmed cases in Sindh (5 deaths) and 16 cases in Islamabad (no deaths), with tertiary care centers seeing mostly complicated or hemorrhagic cases, while primary care saw mild fevers. **Rafique (2015)** found the highest dengue prevalence (57%) in the 21–30 age group, followed by 26% (31–40 years), 11% (41–50 years), and 6.25% (>50 years). Dengue primarily affects young adults and ranges from mild to severe forms. Dengue fever remains endemic in Pakistan, exhibiting year-round transmission with seasonal peaks. However, the unprecedented floods that began in June 2022 created favorable conditions for vector proliferation, resulting in a marked increase in cases compared to previous years. Surveillance data from the National Institute of Health, Islamabad, indicate that between January and September 2022, Pakistan reported 25,932 laboratory-confirmed dengue cases and 62 deaths, with 74% of cases occurring in September alone. Geospatial analysis highlighted uneven provincial distribution: Sindh accounted for 32% of cases (6,888), Punjab including Islamabad Capital Territory 29% (6,255), Khyber Pakhtunkhwa 25% (5,506), and Balochistan 14% (3,128). This outbreak pattern reflects both ongoing endemic transmission and the additional effects of flood-related environmental changes, including expanded mosquito breeding sites and population displacement. These findings underscore the climate-sensitive nature of dengue epidemiology and illustrate how extreme weather events can amplify the burden of arboviral diseases in endemic regions (World Health Organization, 13 October 2022).




Figure 2. Distribution of confirmed dengue cases in Pakistan by province, 1 January to 22 September 2022. Source: World Health Organization (13 October 2022)

Province-District	Year	Duration of the study	Number of Patients	References
KPK				
Peshawar	2010	July to December	38	Usman et al., (2011)
Peshawar, Mardan, Nowshehra, Kohat, Charsadda, Swat	2011	-	139	Ali et al., (2016)
Swat	2013	July to November	6,024	Jehangir et al., (2013)
Swat	2014	July to November	290	Khan et al., (2017)
Swat	2015	July to November	33	
Swat	2016	July to November	45	
Nowshera	2017	July to September	117	
Punjab				
Punjab/2012	2012		40	Zohra et al., (2024)
Punjab/2013	2013		2,647	
Punjab/2014	2014		1,443	
Punjab/2015	2015		4,212	
Punjab/2016	2016		5,053	
Punjab/2017	2017		1,035	
Punjab/2018	2018		961	
Punjab/2019	2019		10,291	
Sindh				
Remedial Centre Karachi	2010	May to October	90	Hasan et al., (2013)
Dengue Control and Prevention Center, East Karachi	2020	September to November	1012	Memon et al., (2024)
Sindh	2000-10		906	Khatak et al., (2023)
Sindh	2011-19		5,416	
Kasmir				

Bagh, Bhimber, Havali, Jehlum Valley, Kotli, Mirpur, Muzaffarabad, Neelum, Rawalakot, Sudhnoti	2016	-	388	Rehman et al., (2018)
Kashmir/2011Govt. Medical College, Hospital, Jammu, and a tertiary care teaching hospital	2015	-	809	Shashi et al., (2016)
	2014	-	78	
	2013	-	6653	
	2012	-	207	
	2011	-	94	

3. Management of Dengue:

3.1. Environmental management

Yousaf (2018) emphasizes that environmental management is a critical component of dengue prevention, particularly in endemic regions. Environmental management involves modifying surroundings to reduce human-vector contact and, consequently, the risk of transmission. Key measures include proper solid waste management, elimination of mosquito breeding sites, improvement of water supply systems, adequate drainage, covering domestic water-storage containers, cleaning flower vases, and recycling old tires. In Pakistan, effective implementation of environmental management by government authorities is essential for controlling dengue. Given the country's ongoing sanitation challenges, the government should prioritize street cleaning, enhanced drainage systems, and removal of stagnant water. At the household level, environmental management is highly effective and should be actively practiced. Additionally, public hygiene campaigns and community cleaning initiatives should be conducted regularly, both prior to and during dengue outbreaks, to minimize transmission risk.

3.2. Biological management of Dengue

Dengue transmission can be effectively reduced through the use of biological control agents that target mosquito populations (Huang et al., 2017). Larvivorous fish, such as mosquito fish, have been widely employed to consume *Aedes* larvae in standing water, while copepods serve as natural predators that feed on mosquito larvae in small water bodies. In addition, certain fungi, including *Beauveria bassiana* and *Metarhizium anisopliae*, have been proposed as potential biocontrol agents due to their ability to infect and kill adult mosquitoes (Benelli et al., 2016). According to Pakistan's national guidelines for dengue control, the introduction of mosquito fish should be limited to small, clean water sources and carried out only after comprehensive surveys of local mosquito breeding sites to ensure their effectiveness and ecological safety. Copepods and fungal agents should also be applied in a targeted manner, focusing on habitats where *Aedes* mosquitoes are most likely to breed (Hammed et al., 2022). Furthermore, innovative approaches such as Wolbachia-based interventions are being actively researched in Pakistan. Wolbachia, a naturally occurring endosymbiotic bacterium, can interfere with the reproduction of mosquitoes and reduce their ability to transmit dengue virus. Early field studies suggest that releasing Wolbachia-infected mosquitoes could significantly decrease dengue transmission over time, making it a promising addition to integrated vector management strategies. Overall, combining these biological methods with environmental management, chemical control, and public health education provides a sustainable and ecologically sound approach to dengue prevention (Yen and Failloux 2020).

3.3. Chemical management

Siddique (2018) highlighted that chemical control is a key component of dengue

prevention. The dengue vector, *Aedes aegypti*, can be managed using larvicides and space sprays. Larviciding is primarily applied to domestic water containers and is generally considered a short-term intervention. Commonly used larvicides for dengue control include pyriproxyfen, *Bacillus thuringiensis* (H-14), and temephos. Space spraying involves dispersing fine insecticide droplets in the air to kill adult mosquitoes and has been the primary method employed by many countries to control dengue fever and dengue hemorrhagic fever. In Pakistan, larvicides are occasionally applied to mosquito breeding sites; however, their use requires careful assessment of breeding areas and strict adherence to safety precautions. Space spraying is typically reserved as an emergency measure during dengue outbreaks to reduce vector populations rapidly.

3.4. Preventive Management

Jahan (2011) emphasizes the importance of prevention and health education in dengue management. Primary care professionals, when equipped with appropriate training, resources, and access to specialist guidance, have the capacity to provide comprehensive care for the majority of patients. General practitioners and community nurses play a pivotal role in promoting health education and hygiene practices within communities (Syed, 2010). Several approaches can be implemented to address the challenge of dengue fever effectively.

3.4.1. Away from contact

The primary preventive measure against dengue involves minimizing contact with infected *Aedes* mosquitoes, which primarily bite during daylight hours. This can be achieved by properly managing waste, improving water storage practices, and eliminating all sources of stagnant water. Additional protective strategies include using household mosquito repellents such as coils, sprays, or nets, and wearing long-sleeved shirts, trousers, and socks to reduce exposure to bites (Idrees, 2012). In Pakistan, particularly in Punjab, the government has been actively promoting dengue prevention through public awareness campaigns aimed at educating communities about the disease and protective measures.

3.4.2. Vaccination

The second approach to dengue control involves vaccination, which is still under development, while the third approach is drug therapy; however, no specific antiviral drugs currently exist to target the dengue virus (Selisko, 2006). Nevertheless, supportive care remains critical and can be lifesaving. Fever is managed with antipyretics such as paracetamol, while joint pain can be relieved using analgesics. Patients with dengue hemorrhagic fever (DHF) or dengue shock syndrome (DSS) require hospitalization. Dehydration can be managed with oral rehydration therapy, and if oral intake is insufficient, intravenous fluid replacement is used to prevent shock. In cases where platelet counts fall below 20,000 or significant bleeding occurs, platelet transfusions are recommended. Medications such as aspirin, ibuprofen, and other non-steroidal anti-inflammatory drugs should be avoided as they may exacerbate bleeding tendencies, and any drugs that reduce platelet counts should also be avoided (Idrees, 2012). To improve dengue surveillance and response, it is essential to establish local laboratories in every union council across the country for accurate case counting, outbreak reporting, and timely implementation of precautionary measures. Additionally, a central research institute for dengue should be established at the district level in each province, linked to all local units, hospitals, laboratories, health centers, and basic health units to facilitate coordinated research and control efforts.

3.4.3. Quick response

The primary objective of dengue research institutes is to provide rapid responses to patients, communities, and government authorities regarding dengue treatment and prevention. To effectively reduce dengue incidence in Pakistan, national dengue control programs must adhere to all WHO-recommended guidelines. Key WHO strategies

include geographically referenced entomological surveillance, active public participation, identification and elimination of mosquito breeding sites, ecological management including pesticide fogging, robust medical examination systems, and community health education. Effective dengue control requires coordinated efforts at local, national, and international levels. Additionally, large-scale, parallel research initiatives should be encouraged across multiple cities in Pakistan. These institutions will also oversee medical evaluations and facilitate the development and implementation of emerging treatments, ensuring that new therapeutic approaches can be safely and efficiently delivered to patients in the future (Zafar, 2013).

3.4.4. Family physicians

The effectiveness of family physicians in utilizing specific communication skills is crucial for improving the physical, mental, and emotional well-being of both patients and their families. It is essential to regularly update the health education system, ensuring that accurate and comprehensive information about dengue fever is widely accessible. Popular media channels, including newspapers and television, should be actively used to disseminate information on a large scale to enhance public awareness and prevention efforts (Ageep, 2006).

3.4.5. Wearing full-sleeve clothes

Dengue-transmitting mosquitoes primarily bite during the daytime. Protection measures include wearing long-sleeved shirts, full-length dresses, and other clothing that covers the limbs, as well as using mosquito repellents, coils, and electric vapor mats during daylight hours. Insecticide-treated nets (ITNs) provide additional protection for vulnerable groups such as young children, pregnant women, and the elderly, as well as for anyone resting during the day. Curtains made of cloth or bamboo can also be treated with insecticides and hung over windows or doorways to repel or kill mosquitoes.

3.4.6. Removal of water containing saucers

Water should be drained from desert or window air coolers when not in use, as well as from tanks, barrels, drums, and buckets. All household items that can hold water, such as plant saucers, should be removed. Stored water containers must be kept covered at all times, and discarded items that can collect water such as bottles, plastic bags, tins, and old tires should be collected and destroyed. Vector control is achieved through a combination of environmental management and chemical interventions (Riaz, 2009).

3.4.7. Proper solid waste disposal

Proper solid waste disposal and improved water storage practices, including covering containers to prevent access by egg-laying female mosquitoes, are promoted through community-based programs (Kay, 2002). The targeted application of insecticides to larval habitats, particularly domestic water storage containers, can effectively prevent mosquito breeding for several weeks; therefore, these treatments should be applied periodically (Hanh, 2009). Recent dengue outbreaks in Pakistan have resulted in substantial morbidity and mortality. Optimal health outcomes depend on timely and accurate diagnosis, along with appropriate treatment. Patient-centered communication plays a critical role in achieving these outcomes, as it provides a comprehensive clinical understanding, facilitating symptom resolution, reduced psychological distress, improved overall health and functional status, effective pain relief, and better management of anxiety.

3.5. Punjab Health Line Project for Dengue Management

Dengue infection imposes a significant burden on both the economic and healthcare systems of Pakistan each year. In response, the Government of Punjab launched the “Punjab Health Line Project for Dengue” to facilitate public awareness of dengue signs and symptoms, provide assistance for suspected cases, and identify areas affected by outbreaks. In early September 2011, the Punjab government temporarily closed schools, colleges, and universities for ten days to conduct intensive mosquito-spraying campaigns. The federal government has invested millions of dollars in various measures to address the outbreak.

Dengue fever has impacted multiple sectors, including the economy, education, and politics, with debates arising among the Punjab government, opposition parties, and media regarding local governance and preparedness. However, the reality is that Punjab, like much of Pakistan, lacks sufficient infrastructure to manage sudden surges of infectious diseases effectively (Tahire, 2012).

4. Conclusion:

Dengue fever remains a persistent public health challenge in Pakistan, with increasing incidence and recurrent outbreaks posing serious clinical and socio-economic burdens. Analysis of regional data highlights significant demographic and age-specific vulnerability, with young adults disproportionately affected. Current management strategies including environmental sanitation, biological and chemical vector control, and preventive measures play an important role in reducing transmission, yet gaps in healthcare provider knowledge and infrastructure limitations compromise optimal outcomes. Family physicians and community health programs are crucial for early detection, patient monitoring, and public education. Integrating innovative approaches such as Wolbachia-based vector control, vaccination programs, and community awareness campaigns with established interventions could significantly reduce dengue transmission. Strengthening national preparedness, improving diagnostic capacity, and promoting continuous research are essential to mitigate future outbreaks and ensure effective clinical and public health management of dengue in Pakistan.

REFERENCES

Ali A, Ahmad H, Idrees M, Zahir F, Ali I. Circulating serotypes of dengue virus and their incursion into non-endemic areas of Pakistan; a serious threat. Virol J. 2016 Aug 26;13(1):144. doi: 10.1186/s12985-016-0603-6. PMID: 27565893; PMCID: PMC5002134.

Begum, F., Das, S., Mukherjee, D., & Ray, U. (2019). Hijacking the host immune cells by dengue virus: molecular interplay of receptors and dengue virus envelope. *Microorganisms*, *7*(9), 323.

Benelli, G., Jeffries, C. L., & Walker, T. (2016). Biological control of mosquito vectors: past, present, and future. *Insects*, *7*(4), 52.

Berube, D. M. (2023). Vectors and reservoirs. In *Pandemics and Resilience: Lessons we should have learned from Zika* (pp. 265-302). Cham: Springer International Publishing.

Bhugra, A. (2020). *Correlation of Circulating Levels of Free Secreted Dengue Nonstructural 1 (sNS1) Antigen with Severity of Disease in Adult Patients Admitted with Dengue* (Doctoral dissertation, Rajiv Gandhi University of Health Sciences (India)).

Dehghani, R., & Kassiri, H. (2021). A review on epidemiology of dengue viral infection as an emerging disease. *Research Journal of Pharmacy and Technology*, *14*(4), 2296-2301.

Dowd, K. A., Sirohi, D., Speer, S. D., VanBlargan, L. A., Chen, R. E., Mukherjee, S., ... & Pierson, T. C. (2023). prM-reactive antibodies reveal a role for partially mature virions in dengue virus pathogenesis. *Proceedings of the National Academy of Sciences*, *120*(3), e2218899120.

Ferreira-de-Lima, V. H., & Lima-Camara, T. N. (2018). Natural vertical transmission of dengue virus in Aedes aegypti and Aedes albopictus: a systematic review. *Parasites & vectors*, *11*, 1-8.

Hamed, A. M., El-Sherbini, M. S., & Abdeltawab, M. S. (2022). Eco-friendly mosquito-control strategies: advantages and disadvantages. *Egyptian Academic Journal of Biological Sciences, E. Medical Entomology & Parasitology*, *14*(1), 17-31.

Hasan SR, Riaz M, Jafri FA. Characteristics and outcome of dengue infection; clinical perspective from a secondary care hospital of Karachi. Pak J Med Sci. 2013 Jan;29(1):115-8. doi: 10.12669/pjms.291.2742. PMID: 24353520; PMCID: PMC3809171.

Htun, T. P., Xiong, Z., & Pang, J. (2021). Clinical signs and symptoms associated with WHO severe dengue classification: a systematic review and meta-analysis. *Emerging microbes & infections*, *10*(1), 1116-1128.

Jehangir Khan, Asar Khan. Incidence of dengue in 2013: Dengue outbreak in District Swat, Khyber Pakhtunkhwa, Pakistan. Int. J. Fauna Biol. Stud. 2015;2(1):01-07.

KARISA, J. K. (2018). *URBAN ECOLOGY OF CULEX QUINQUEFASCIATUS AND AEDES AEGYPTI MOSQUITOES AND TRANSMISSION DYNAMICS OF ARBOVIRUSES IN COASTAL KENYA* (Doctoral dissertation, Pwani University).

KARISA, J. K. (2018). *URBAN ECOLOGY OF CULEX QUINQUEFASCIATUS AND AEDES AEGYPTI MOSQUITOES AND TRANSMISSION DYNAMICS OF ARBOVIRUSES IN COASTAL KENYA* (Doctoral dissertation, Pwani University).

Khan W, Khan BA, Khan Z, Abid ur Rehman, Akbar M, Khan IA. Dengue fever; the clinical pattern and mortality in epidemic and post epidemic years in Swat. Professional Med J 2017;24(10):1466-1470. DOI:10.17957/TPMJ/17.3947

Khattak, A., Khan, S., Ali, I., Gul, A., Khabir, M. N., Javed, B., Ayesha, Adnan, M., Khan, S. N., & Attaullah, S. (2023). Burden and distribution of dengue infection in Pakistan (2000-19): a review. *Brazilian Journal of Biology*, *84*, e267982. https://doi.org/10.1590/1519-6984.267982

Khattak, A., Khan, S., Ali, I., Gul, A., Khabir, M. N., Javed, B., Ayesha, Adnan, M., Khan, S. N., & Attaullah, S. (2023). Burden and distribution of dengue infection in Pakistan (2000-19): a review. Brazilian Journal of Biology, 84, e267982. https://doi.org/10.1590/1519-6984.267982

Kok, B. H., Lim, H. T., Lim, C. P., Lai, N. S., Leow, C. Y., & Leow, C. H. (2023). Dengue virus infection–a review of pathogenesis, vaccines, diagnosis and therapy. *Virus research*, *324*, 199018.

Krishna, G., Sreedevi, S., & Thrimothi, D. (2024). Dengue Virus Infection: Etiology, Epidemiology, Pathogenesis, Diagnosis, and Prevention. In *Viral Infectious Diseases Annual Volume 2024*. IntechOpen.

Li, Q., & Kang, C. (2022). Structures and dynamics of dengue virus nonstructural membrane proteins. *Membranes*, *12*(2), 231.

Lin, D. C. D., Weng, S. C., Tsao, P. N., Chu, J. J. H., & Shiao, S. H. (2023). Co-infection of dengue and Zika viruses mutually enhances viral replication in the mosquito Aedes aegypti. *Parasites & Vectors*, *16*(1), 160.

Mahajan, A. (2018). *A Study on Hypocalcemia and Its Potential Value as a Biochemical Marker in Predicting Severity of Dengue Infection* (Doctoral dissertation, Rajiv Gandhi University of Health Sciences (India)).

Mboera, L. E., Mweya, C. N., Rumisha, S. F., Tungu, P. K., Stanley, G., Makange, M. R., ... & Oriyo, N. M. (2016). The risk of dengue virus transmission in Dar es Salaam, Tanzania during an epidemic period of 2014. *PLoS neglected tropical diseases*, *10*(1), e0004313.

Mehmood, A., Khan, F. K., Chaudhry, A., Hussain, Z., Laghari, M. A., Shah, I., ... & Ikram, A. (2021). Risk factors associated with a dengue fever outbreak in Islamabad, Pakistan: Case-control study. *JMIR Public Health and Surveillance*, *7*(12), e27266.

Memon A, Tahir S, Tahir SA. Dengue Outbreak, a growing endemic – an overview of dengue cases in Karachi. Professional Med J 2024; 31(06):856-861. https://doi.org/10.29309/TPMJ/2024.31.06.8133

Muller, D. A., Depelsenaire, A. C., & Young, P. R. (2017). Clinical and laboratory diagnosis of dengue virus infection. *The Journal of infectious diseases*, *215*(suppl_2), S89-S95.

Murugesan, A., & Manoharan, M. (2020). Dengue virus. In *Emerging and reemerging viral pathogens* (pp. 281-359). Academic Press.

Pourzangiabadi, M., Najafi, H., Fallah, A., Goudarzi, A., & Pouladi, I. (2024). Dengue virus: Etiology, epidemiology, pathobiology, and developments in diagnosis and control–A comprehensive review. *Infection, Genetics and Evolution*, 105710.

Rehman SN- ur-, Hafeez U, Gafoor T, Rana A. Epidemiological Profile of Dengue Fever Cases in Azad Jammu & Kashmir, Pakistan. Int J Pathol [Internet]. 2018 Nov. 21 [cited 2025 Jul. 20];:55-9.

Roy, S. K., & Bhattacharjee, S. (2021). Dengue virus: epidemiology, biology, and disease aetiology. *Canadian journal of microbiology*, *67*(10), 687-702.

Shashi, S. S., Sharma Monika, Kumar, G. R., & Sorabh, S. S. (2016). Sero- Epidemiological trends of Dengue Fever in Jammu Province of JandK State. International Journal of Medical Research & Health Sciences, 5(8), 1–6.

Songprakhon, P., Thaingtamtanha, T., Limjindaporn, T., Puttikhunt, C., Srisawat, C., Luangaram, P., ... & Noisakran, S. (2020). Peptides targeting dengue viral nonstructural protein 1 inhibit dengue virus production. *Scientific Reports*, *10*(1), 12933.

Sun, Y. T., & Varani, G. (2022). Structure of the dengue virus RNA promoter. *RNA*, *28*(9), 1210-1223.

Ten Bosch, Q. A., Clapham, H. E., Lambrechts, L., Duong, V., Buchy, P., Althouse, B. M., ... & Perkins, T. A. (2018). Contributions from the silent majority dominate dengue virus transmission. *PLoS pathogens*, *14*(5), e1006965.

USMAN, M., JAMAL, S., TABASSUM, S. and ZAFRAN, M., 2011. Outcome of dengue fever in adults. Gomal Journal of Medical Sciences., vol. 9, no. 1, pp. 70-73.

Venkatesan, P. (2024). Global upsurge in dengue in 2024. *The Lancet Infectious Diseases*, *24*(10), e620.

World Health Organization (13 October 2022). Disease Outbreak News; Dengue - Pakistan. Available at: https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON414

Yen, P. S., & Failloux, A. B. (2020). A review: Wolbachia-based population replacement for mosquito control shares common points with genetically modified control approaches. *Pathogens*, *9*(5), 404.

Zohra, T., Din, M., Ikram, A. *et al.* Demographic and clinical features of dengue fever infection in Pakistan: a cross-sectional epidemiological study. *Trop Dis Travel Med Vaccines* 10, 11 (2024). https://doi.org/10.1186/s40794-024-00221-4