The Research of Medical Science Review
ISSN: 3007-1208 & 3007-1216 Volume 4, Issue 3, 2026

# BLOWING AWAY THE COMPETITION: THE CRITICAL INSIGHT ABOUT SECRET ECOLOGICAL POWER OF PUFFERFISH

**Shahid Mahmood[*1], Rubab Fatima[2], Syeda Marjan Zahra Naqvi[3], Arooba Inayat[4], Aleeza Parvez[5], Laiba Shama[6], Pakeeza Shahzadi[7]**

[*1,2,3,4,5,6,7]*Department of Zoology, University of Gujrat, Gujrat 50700, Pakistan*

[*1]shahid.mahmood@uog.edu.pk

DOI: https://doi.org/10.5281/zenodo.19329376

**Keywords**
*Puffer fish, Habitat distribution, Feeding ecology, trophic interaction, Toxicity*

**Article History**
*Received: 28 January 2026*
*Accepted: 15 March 2026*
*Published: 30 March 2026*

**Copyright@Author**
**Corresponding Author: ***
**Shahid Mahmood**

**Abstract**
*Millions of organisms live in the water, but only a small number are truly unique. The puffer fish is one of these odd creatures that, despite being the second most poisonous vertebrate, has distinctive qualities many people find fascinating. Predators find it more difficult to consume puffer fish, also called blowfish, swellfish, and globefish, because they can inflate themselves with water or air when attacked. These fishes are classified as tetrodotoxic. The three families of puffer fish, called Fugu in Japan, are Tetraodontidae, Triodontid, and Diodontidae. The Tetraodontidae family includes species found in marine, brackish, and freshwater environments. The Tetraodontidae refers to four teeth used to break the shells of molluscs and crustaceans. These teeth are arranged into two upper plates and two lower plates.*

IEER
Institute for Excellence in Education & Research

## INTRODUCTION

The Tetraodontidae family includes 30 genera and 200 species and lives in freshwater and shallow, warm tropical oceans worldwide. A massive blunt head, jaws shaped into a beak with strong heavy teeth including two above and two below, gill openings that look like simple slits in front of the pectoral fins, and the absence of pelvic fins are some physical characteristics of this family. Tetraodontiforms belong to the Tetraodontidae family and represent a distinct monophyletic group within Teleostei. Because of its distinctive physical characteristics, ichthyologists and biologists have long been interested in this group. Tetraodontidae is the only family of tetraodontiforms that lives in freshwater. One of the main challenges aquatic creatures face when expanding their range is crossing the freshwater-sea boundary, since it involves bridging difficult gaps between different environments. In this sense, this family of tetraodontiforms is considered the most unique and intriguing. (Nelson *et al.*, 2016).

That every fish's reproductive habits are closely connected to its evolutionary background. Information on the various reproductive strategies of Tetraodontid fishes can help explain their evolutionary history, including how the Tetraodontidae adapted to freshwater environments. The distinctions between freshwater and brackish water puffer fishes' reproductive behaviors have not been well studied. There are only 10 documented cases of captive breeding of freshwater and brackish water puffer fish. One example is the African *Tetraodon schoutedeni*, which spawns by swimming alongside one or more males.

The females bite and hold onto their abdomens as they lay demersal eggs. Before discussing the distinctive traits of puffer fish that likely help them succeed in their invasion, some background information is provided. Except when a specific species is mentioned, in which

case its scientific name is provided, both. In this study, they are referred to as puffer fish. A complete knowledge of these incidents is necessary to build this baseline database of occurrences, assess temporal and regional changes in threat, and develop more effective outreach and preventive initiatives for the wider Mediterranean region.(Ulman *et al.*, 2024)

In 2024 conducted a study that explored various aspects of the subject. Their findings provided valuable insights that helped shape further research and understanding. Marine Protected regions (MPAs) are legally protected areas that stretch from the intertidal zone to the deep ocean. They are created to conserve endangered species and ecosystems by maintaining biodiversity and reducing the risk of extinction for certain species. The restoration of environmental integrity should be possible through MPAs by increasing marine output, including fish and invertebrate species. Research has shown that fish serve as an important tool to monitor the impact of human activity on marine ecosystems and to assess the environmental quality of coastal areas.(Marinsek *et al.*, 2024)

*Sphoeroides testudineus* is a puffer fish found widely in the Eastern Atlantic and is common along the Brazilian coast. They are vital to maintaining the trophic balance in bays and estuaries, where they like to reside. They mostly eat gastropods, bivalves, and crustaceans. Because *S. Testudineus* fish can tolerate a wide range of environmental conditions and have been used as sentinel in multi-level biomarker studies (Gonçalves, Andriolo, Pinto, & Duarte, 2020). The performance of fish biomarkers depends on the sensitivity of the species being studied, the biomarker selection, and the conditions of sampling and analysis. These difficulties can be reduced by careful experiment planning and execution, along with a critical evaluation of the results. Organs like the liver and gills of fish are used as targets in ecotoxicological studies because they show how contaminants might affect gas exchange and detoxification. The gills are an organ that is in direct connection with the environment and with the materials present in the water. They are important for both breathing and ion exchange. The gills' histological changes are typically linked to acute contamination effects such as hypertrophy, necrotic patches, lamellar fusion, and elevation of the lamellar epithelium. Changes in gill shape may be closely linked to problems with gas exchange and osmoregulation because they increase the distance between epithelial cells and blood capillaries. This can compromise osmoregulation when hazardous chemicals are present. Steatosis, circulation issues, melanomacrophage centers, and necrotic areas are some of the alterations observed. (Suchana *et al.*, 2021)

China and Japan, puffer fish are considered a valuable aquatic product and have been eaten for a long time because they contain high protein and have a tasty flavor. Despite their great flavor and high market value, many puffer fish contain unhealthy levels of tetrodotoxin (TTX) in their ovary, liver, kidney, eyes, and blood (Wu & Chen, 1981). Ingestion or incorrect handling can cause poisoning or even death. It is essential to filter the edible puffer fish species to avoid poisoning from eating hazardous puffer fish. Identification and taxonomy of puffer fish provide solutions to the problems described above. There have been many reports on puffer fish molecular taxonomy so far. These studies mainly focused on a few genera or species and did not include a comprehensive taxonomic study of the species across the entire order Tetraodontiformes. There is an urgent need for research on the classification and identification of puffer fish using DNA barcoding to clarify the cryptic species and synonyms of this order.(Liu *et al.*, 2024).

The Suez Canal, transportation, connecting straits, global warming, and climate change all contribute to the growth and spread of invasive foreign marine creatures in the Mediterranean Sea. Because of global warming and climate change, the Mediterranean Sea is turning into a tropical sea. As invasive alien marine species increase, local marine biodiversity and the Mediterranean Basin's sustainable seafood market face serious risk. The literature covers many important subjects, including attacks by invasive alien species that Mediterranean countries face. Physically identifying and tracking alien species in invasive alien studies is difficult, but researchers make several assumptions about the effects of alien species on

the waters and efforts to control them (Kaplan, Yildirim, & Yildirim, 2022).

**1. Taxonomy, Morphology and Distribution**

Puffer fish provide important information for understanding these fish. Taxonomy helps classify puffer fish within the animal kingdom morphology studies their physical characteristics, such as body shape and skin texture. Distribution refers to where puffer fish are found in oceans and coastal areas. This knowledge supports research and conservation efforts related to these species. Puffer fish belong to the family Tetraodontidae and are known for their ability to inflate their bodies and for having complex toxicity mechanisms. Because these species have specialised roles in benthic food webs and are often used as markers of reef health, understanding their taxonomy, morphology, and geographic distribution is essential. It has been established that creating algorithms for the individual identification of animals from images is ecologically important. Most algorithms are trained separately for each species, which limits their widespread use because data curation and model maintenance are expensive (Otarashvili et *al.,* 2024).

The Tetraodontidae family includes puffer fish and stands out as one of the most physically specialised and evolutionarily unique families within the Tetraodontiformes order. Puffer fish are a unique group of fishes known for their protective inflating mechanism and the poison tetrodotoxin. Puffer fish belong to the category Eukaryotes, which includes organisms with cells that have specialised organelles and a membrane-bound nucleus. They belong to the kingdom Animalia within this domain because they are heterotrophic, multicellular organisms that have complex tissue structures and the ability to move (Nelson et *al.,* 2016).

At the phylum level, puffer fish are part of the Chordata group. This group is defined by features such as a post-anal tail at some point in their development, pharyngeal slits, a dorsal hollow nerve cord, and a notochord. During the development of the embryo, these characteristics are especially noticeable. (Helfman et *al.,* 2009). The largest group of vertebrates, the Actinopterygii or ray-finned fishes, includes species such as the pufferfish. They are distinguished by the fact that their fins are supported by bony rays rather than fleshy lobes (Helfman et al., 2009). Puffer fish belong to the Tetraodontiformes order of the Actinopterygii, which is known for its highly specialized body structure.This order of fishes often has fused teeth, changed scales, and fewer skeletal features.Their distinct eating habits and protective tendencies are a result of these adaptations.The order includes various odd fish families such as sunfish, boxfish, and triggerfish, but puffer fish are known for their exceptional ability to inflate their bodies when they sense danger (Froese & Pauly, 2024). There are two main reasons current methods of monitoring marine species are inadequate. First, existing approaches sometimes overlook the similarities in appearance between related species, requiring large datasets for each specific species something uncommon or cryptic puffer fish often lack (Otarashvili et al., 2024).

Second, the fine-scale spatial resolution needed to link species occurrence with specific microhabitats is often missing from regular monitoring. Coarse-scale measurements may miss important early warning indicators of environmental change because species are very sensitive and respond to small differences in their microhabitats, as shown in studies of different benthic assemblages. (Lasinio et *al.,* 2017).

**1.1. Multi-Species Animal Re-Identification**

Involves recognizing and matching individual animals across different species based on their unique features. This process helps in tracking wildlife, studying animal behavior, and supporting conservation efforts while ensuring accurate identification regardless of species differences. Recent advancements have shown that training a single embedding network for multiple species, including whales and primates, can improve performance (Otarashvili et al., 2024).

This approach is particularly relevant for pufferfish taxonomy, where shared morphological traits play an important role e.g. Variations in body shape and fin placement across the Tetraodontidae family can help

improve the identification of rare species when there is limited training data. By focusing on identifying animals individually, this method helps in studying population dynamics and morphology over time. (Otarashvili et al., 2024). Puffer fish include about 28 genera and more than 190 species. They live in different environments, such as shallow tropical coral reefs, temperate open ocean areas, and freshwater rivers in South America, Africa, and Southeast Asia. (Zhang et al., 2025).

The evolutionary trajectory of this group includes the reduction of skeletal elements, the fusion of teeth into a beak-like structure, and the development of high elasticity in the ventral skin to facilitate inflation. (Zhang et al., 2025). Pufferfish in reef environments share similar characteristics, often appearing cryptic or camouflaged. Research into detecting small and rare wildlife in aerial imagery has led to the development of frameworks like RareSpot, which integrates multi-scale consistency learning to reduce feature loss related to scale. (Zhang et al., 2025)

For instance, color patterns, which are often primary diagnostic features, can vary greatly depending on life stage, sexual dimorphism, and environmental background matching. The phylogenetic relationships between marine and freshwater clades are not fully resolved, which complicates our understanding of how freshwater invasion events occurred historically (Otarashvili et *al.,* 2024). Pufferfish from the family Tetraodontidae live in many areas around the world. They can be found in marine, brackish, and freshwater environments in tropical and subtropical regions.

### 1.2. Morphological Taxonomy of Fish (Classical)

In the past, taxonomists relied on three main character types to classify fish: the structure of the nasal apparatus, the presence/absence of a lateral line, and the dermal spine distribution (spinules). For example, the nasal organ (with a single tentacle, bifid structure, or an open pit) helped biologists distinguish the genera Arothron and Canthi aster through the application of dichotomous keys. Why use a classical taxonomic approach? The classic approach is both accessible and able to identify species in the field without expensive lab equipment (Croze et al., 2019).

However, one significant limitation is that the features upon which the classical method relies are subject to ontogenetic change; i.e., juvenile pufferfish display nasal structures/spinal patterns that differ greatly from those of adults, and, historically, taxonomists have synonymized different life stages of the same fish species as two separate species. Also, preservation of delicate dermal spines also obscures important diagnostic information. (Lasinio et al., 2017).

### 1.3. Ecological Distribution and Toxicology

The other area of researched relationships is the relationship between the ecological distribution of pufferfish and the distribution of toxic compounds. The study is focused on how the amounts of tetrodotoxin (TTX) and saxitoxin that are sequestered in different tissues (liver, ovaries, skin) and geographical areas (Otarashvili et al., 2024). The icebergs of ecological distribution state that the toxicity of pufferfish acts as a selective pressure for the distribution of pufferfish and how behaviourally and morphologically (i.e., aposematism) individuals are different. Research shows that there is often an association between the number of toxic pufferfish and the presence of TTX-producing bacteria within the local food web (Zhang et al., 2025).

## 2. Feeding Ecology and Behavioral Ecology of Pufferfish

The family of pufferfish (known scientifically as Tetraodontidae) is comprised of teleost (bony) fishes and includes representatives that inhabit all types of aquatic ecosystems ranging from tropical to temperate marine and freshwater habitats. One reason for their ecological success is that they have adapted morphologically and behaviorally to their surroundings. These adaptations include strong beak-shaped dentition, which enables them to consume the hard bodied prey. The fact that the teeth of pufferfish are fused together in the form of a beak results in their ability to consume durophagically (the consumption of mollusks, crustaceans, and other kinds of hard-bodied prey) makes them different from most other

families of teleost fishes and clearly demonstrates their role in a different trophic level than the other families of teleost (Evans et al., 2023; Tyler, 1980). The morphological innovation associated with the beak dentition lends considerable support for much of the feeding ecology associated with pufferfishes and enables them to exploit and utilize a wide variety of ecological niches, including coral reefs, estuary habitats, and freshwater habitats.

### 2.1. Feeding ecology

There are many unique characteristics of pufferfish which ultimately lead to their dietary differences depending on the specific habitat they occupy, the species, and their respective ontogenetic stage of life. (Renjithkumar et al., 2020).

Renjithkumar, & Ranjeet evaluated the dietary characteristics of *Carinotetradon travancoricus*, a species of freshwater endemic pufferfish from the Western Ghats of India. The authors reported that the diet of juvenile C. *travancoricus* is primarily omnivorous and that the diet of adult C. *travancoricus* is primarily insectivorous and crustaceanivorous. Feeding behaviour is also affected by seasonality, with the intensity of feeding occurring prior to the monsoon (higher levels of feeding), and as spawning approaches, the diet shifts to become more animal based (i.e., more insects and/or crustaceans), indicating that C. *travancoricus* exhibits a high degree of flexibility in their respective diets, as influenced by life stages and environmental conditions (Renjithkumar et al., 2020). To better understand feeding plasticity, the following diagram presents the ontogenetic dietary shifts of *Carinotetraodon travancoricus* in relation to developmental stages and habitat variability.




**Fig. 01: Dietary Shift of *Carinotetradon travancoricus* (Flexible feeding Adaptations Based on Life Stages and Environment)**

### 2.2. Development and seasonal variability of appetite

Puffers' feeding patterns are determined by both their age and the environment in which they live. For example, the diet of *Colomesus asellus* in the upper-middle Tocantins River, Brazil, was primarily made up of benthic insects (mostly immature mayflies), and it did not vary systematically with either geographic location or size. Hence, it can be inferred that trophic roles in this population's feeding ecology will not vary as much between seasons and developmental stages (Bartolette et al., 2018).

However, some species exhibit seasonal variation with respect to foraging behaviour related to their life cycle. For example, *Carinotetradon travancoricus* exhibited greater intensity of foraging on animal food items during the spawning cycle than at other times of year (Renjithkumar et al., 2020). Ultimately, while there are systems where fish diets remain fairly constant across freshwaters, unique characteristics such as reproduction will affect feeding behaviors differently in other environments.

### 2.3. Intermediate Trophic Position and Energy Transfer through Trophic Relations

Finally, in ecosystems, *C. psittacus* serves a critical role through its position in the food web as both predator of invertebrates living near the water's bottom and prey for larger fish species. For instance, *C. psittacus* in mangrove areas have been shown to consume large numbers of barnacles and crabs; thus, there is a high energy transfer through these fisheries across estuarine ecosystems. Based on the analysis of stomach contents of *C. psittacus* in these areas, it was found that approximately 6.2% of their total body weight was consumed daily in barnacles or 178.7 grams/ha/day in brachyuran crabs, further illustrating their role as an efficient conduit of energy through these fisheries (Krumme et al., 1997).

### 2.4. Behavioural Ecology: Foraging Behaviour and Social Interactions

Feeding behaviour may also influence social dynamics among pufferfish, as evidenced by research in Behavioural ecology. For example, in juvenile tiger puffer (*Takifugu rubripes*), restricting food intake leads to an increase in aggression; increased amount of activity; and increases in body contact with other fish while feeding. Brain monoamine levels have been shown to be markedly altered due to reduced availability of food and therefore intensification of intraspecific competition. Elevated rates of dopamine and reduced levels of 5-HIAA and MAOA indicate that restricted food availability may influence aggression through changes to neurological processes (Xie et al., 2021). Aggressive behavior will have ecological consequences in terms of territorial behaviour, dominance hierarchies, and resource access in natural populations, particularly in environments where food is limited or patchily distributed. The behaviour exhibited by fish will significantly affect their fitness and, subsequently, community structure.

### 2.5. Defensive Behaviour and its Ecological Context

Pufferfish are well known for their inflation behaviour, which is a rapid defensive behaviour used by pufferfish to defend themselves from predators. Inflating results in increased size and corresponding reduced chances of predation as a result of gulping water or air into their elastic stomachs. The anatomy, skin elasticity and musculature, and neural control of the inflation behaviour in *Takifugu obscures* have all demonstrated their role in assisting the function of the pufferfish in their defense mechanism (i.e., a functional role in survival).

In addition to using physical means to defend themselves, many species of pufferfish are able to produce tetrodotoxin (TTX), which is derived primarily from their food supply or their microbial flora and is used as a chemical defense. Studies have shown that TTX can reduce the number of attacks from predators, and in juvenile puffer abundance, TTX may reduce stress and agonistic interactions between conspecifics (Ferreira et al., 2022). Thus, it appears that the roles of chemical and physical defense, feeding ecology, and Behavioural ecology are related in pufferfish.

### 2.6. Ecological Factors Affecting the Management of Puffer Fish

Puffer fish are among some of the earliest species to inhabit marine environments. Their discovery is closely linked with discoveries of; the first chlorine dioxide (ClO2) reactions and ignition of electricity by using electrolysis. Marine puffer fish are common in coral reef and seagrass habitats; both of which frequently have high prevalence of mollusks, urchins, and crustaceans. Many studies have demonstrated that durophagy feeding is a common phenomenon across all puffer fish (Evans et al., 2023).

Conversely, freshwater species of puffer fish, including both *Colomesus asellus* and *Carinotetradon travancoricus*, are most often feeding on benthic insects and smaller invertebrates. The feeding habits of these freshwater species, therefore reflect the types of lentic or lotic ecological systems they inhabit (Bartolette et al., 2018; Renjithkumar et al., 2020).

## 3. Computational Modelling of the Relationship Between Inflation and Tetrodotoxin (TTX) in an Ecologically Functioning Puffer Fish Population

The ecological importance of a puffer fish comes largely from the fact that they possess various specialized behavioral defense strategies that allow them to be able to exist in specific ecological niches, even though many individual pufferfish species are unable to swim quickly. Inflation response is one of the more common defense strategies among all animal species and is often used as a predator avoidance strategy. Up until recently, it has been accepted that inflation response represents a 'breath-holding' response whereby final gill oxygen absorption occurs during inflation process and, simultaneously, the dermal respiration occurs to maintain energy to complete the inflation response (McGee et al., 2014). The two main defence strategies employed by all pufferfish are body inflation, where they expand their bodies to an inflated size that is typically several times their normal size, and use of Tetrodotoxin (TTX) as a chemical means of defending themselves from predation.

Strong esophageal and pyloric sphincters are believed to facilitate the inflation response. The anatomical location of the sphincters (between the esophagus and stomach, and between the stomach and intestine) has been well-documented early on in the anatomy and highlighted as an integral factor in preventing the emptying of the pufferfish stomach with respect to the water that was ingested by the fish (McGee et al., 2014).

The most potent neurotoxin known, tetrodotoxin (TTX), was originally thought to be unique to pufferfish. Still, the determination of its structural complexity has identified TTX in other marine animals, such as the toxic goby (Yonge ichthys criniere), xanthid crabs, blue annulated devilfish, and flatworms as well as amphibians and terrestrial annelids. Since then, a number of TTX analogs including deoxy TTXs, dideoxides, and 5,6,11-trideoxyTTX have been identified in TTX-containing organisms (Asano et al., 2023).

When a pufferfish inflates, it is much more difficult for a predator to swallow and poses a physical barrier to predators that have a restricted biting capacity. In contrast, TTX serves as a lethal deterrent to those predators that manage to avoid being impeded by the morphological defense of the pufferfish. Accordingly, these two methods of deterrence work together to reduce predator-related mortality of pufferfish and, in turn, influence the population dynamics of both pufferfish and their predators. Given the rapid changes occurring in marine ecosystems as a consequence of anthropogenic climate change, it is essential to understand the stability and variability of these defensive mechanisms (Smith et al., 2016).

### 3.1. Environmental Factors of Marine Toxicity

Although environmental (abiotic) influences play a major role in the synthesis and accumulation of toxins in marine organisms, historically, establishing causal relationships has been challenging. Recently, with advancements in deep learning, it is now possible to better predict the presence of toxins by using meteorological and hydrological data.

In research conducted by (Schoppema et al., 2025) on the key drivers of tetrodotoxin (TTX) contamination in bivalve mollusks, explainable AI (XAI) was used to identify solar radiation, water temperature, and chloride concentration as critical predictors; therefore, the definitive environmental factors supporting such correlations were established in their research.
Findings from (Schoppema et al., 2025) will be particularly relevant to puffer fish, given that TTX is often accumulated via the food chain or from symbiotic bacteria that are highly sensitive to many of these same environmental variables. This study also represents a substantial improvement over the biological sampling methods, which capture toxins in a static manner, by using a dynamic, temporal framework to define fluctuations in toxicity.

### 3.2. Adversarial Dynamics and Resilience

Biological systems and computational systems both utilize defensive strategies, either evolutionarily (biological systems) or organically/systematically developed (computational systems), to withstand the effects of "attackers" (adversarial entities); thus adversarial robustness refers to how well a given system resists adversarial-based failure when provided with an input generated by an attack on the method of operation of a given system, such as Fast Gradient Sign Method (FGSM) and Carlini-Wagner (CW), as applied to neural networks (Sarkar et al., 2024).
In ecology, the process of adaptation to changing conditions relies upon a predator, as an example of an adversary, to develop a strategy to optimally overcome the defenses of prey while simultaneously expending as few metabolic resources as necessary. In addition, the manner in which sparse and intentionally applied modifications (attacks) destabilize predictive models has been examined via multivariate time-series forecasts (Liu et al., 2022). which relates directly to evolutionary arms races as the predator adopts a novel behavioral mechanism for bypassing the defensive tactics of prey (i.e. pufferfish), resulting in the necessary evolution of punterfish having many strong defenses in order that their defenses would be effective when under potentially noisy conditions and/or disturbed environmental signals (Liu et al., 2022).

**Table 01: Summary of key ecological features of the selected pufferfishes.**

| Species | Habitat | Distribution | Diet | Defense | Key Feature | Ref. |
|---|---|---|---|---|---|---|
| *Colomesus asellus* | Freshwater | Tocantins River, Brazil | Benthic insects | Inflation; TTX | Elastic skin; beak-like teeth | ((Nelson *et al.*, 2016) |
| *Colomesus psittacus* | Brackish (mangrove) | N. Brazilian coast | Barnacles, crabs | Inflation; TTX | Important mangrove energy link | (Barletta & Blaber et al., 2007) |
| *Takifugu rubripes* | Marine | Japan, China | – | Inflation; TTX | Genomic model species | (Noguchi & Arakawa, 2008). |
| *Sphoeroides testudineus* | Marine/Brackish | Atlantic; Brazil | Mollusks, crustaceans | Inflation; TTX | Environmental sentinel species | (Ferreira et al., 2010) |
| *Takifugu obscurus* | Marine/Brackish | East Asia | – | Elastic stomach; TTX | Specialized inflation anatomy | (Yang & Chen et al., 2003) |
| *Tetraodon schoutedeni* | Freshwater | Africa | – | Inflation; TTX | Demersal egg laying | (FAO, 2023) |
| *Takifugu alboplumbes* | Marine | Pacific coast Japan | – | High TTX (ovary/liver) | Toxin via diet/microbes | (Noguchi & Arakawa et al., 2008) |

### 3.3. Defense Deployment Optimization

An important aspect of ecological defense is the allocation of mass usage-based energy budgets to each of the predator's different forms of defense, whereby, regardless of the number of total predatory defense mechanisms available, an organism cannot maximize all predatory defense mechanisms at once, but must optimize its use based upon its particular energy availability-based budget constraints. In assessing predatory defense deployment methods and strategies available to them in modern contexts, (Ge et al., 2019).

### 3.4. Eco-Defense Simulation Module (EDSM)

For the Eco-Defense Simulation Module (EDSM), we plan to develop a computational framework to study the effectiveness of defensive mechanisms of puffer fish in relation to the surrounding environment. This framework is based on the prediction of biological survival through the analysis of the environment (resources present and temperature) and the defense mechanisms of the puffer fish (inflation and toxicity).

- **Ecological Feature Extraction:** This module collects and processes abiotic data to estimate the potential toxicity of an environment. Following the methodology developed for TTX in mollusks, we propose an LSTM approach to model time series data of parameters such as water temperature, global radiation, and salinity (Schoppema et al. 2025). It is hypothesized that as water temperature and radiation increase, the growth of TTX-producing bacteria accelerates, and thus the chemical defense of the puffer fish increases.
- **Adversarial Interaction Modeling:** This module approaches the simulation of predator-prey interactions as an optimization problem. The predator is an adversarial agent whose goal is to minimize the "cost" of the predation (i.e., the time required to capture and the risk of exposure to the puffer fish's toxin), while the puffer fish is a defender whose goal is to maximize the "cost" of the predator. (Sarkar et al., 2024).

## 4 Reproductive Ecology and Ecological Function in Puffer Fish

Pufferfish make up an important part of coastal and estuarine biodiversity, acting as both a food source for top-level consumers and as a consumer's main food source. Their ecological importance has been frequently diminished by their poisonousness; however, they provide insight into the complex and intricate relationship between reproductive ecology and environmental dependability, both of which play a major role in community structure. Within this context, the genus Takifugu has emerged as a model organism for genomic work, but applying the findings of these studies to macroecology is a complete research gap that must be filled moving forward. The continuance or failure of pufferfish populations is dependent primarily upon the timing of reproductive events and their ability to disperse through larval transport. Both biological rhythms and physical-oceanographic forcing impacts all of the above.

Current ecological methods are often not linking molecular biology to ecosystem dynamics outside of using the genome of *Takifugu rubripes* which is now known. Traditionally, ecological monitoring methods have used visual census or catch data and are very labour intensive, and do not have the resolution necessary to monitor rapid reproductive events or larval dispersal events over time. Traditional methods cannot also provide enough data to understand the degree of "self-organized criticality" in population dynamics; however genomic data would provide insights of that type. Secondly existing species interaction models generally rely on static pair wise interaction matrices which do not help characterize the temporal nature of ecological interactions. Failure to account for time-dependent sign changes in species interactions will lead to misinterpretation of the importance of puffer fish fluctuating roles as they transition from larvae to adult (Aguilar et al., 2025). Thus our understanding of the ecological role of puffer fish is still not complete without considering time dependent sign changes in puffer fish vs puffer fish interactions. The fractal nature of many biological signals, from the series of heartbeats to DNA sequences, is evident through the analysis of groups of data. Selvam has conducted extensive research on the

*Takifugu rubripes* genome (the puffer fish), and it is well established as an example of this pattern of fractal variation in biological data series. He has also determined that the frequency distribution of DNA bases in the *Takifugu rubripes* genome follows a universal inverse power law (Selvam, 2007). This is an indication of space-time long-range correlations or self-organising criticality, suggesting that the non-coding regions of this genome are not "junk," but serve an important function in the integration of systems (Selvam, 2008).

While this literature is a strong statistical illustration of the *Takifugu rubripes* genome, it deals primarily with the internal physical properties of the DNA sequence. We then go further to propose a hypothesis regarding the connection of these fractal genomic signatures to the adaptive ability of the organism to the fractal nature of fluctuations in the surrounding environment, which is only partially addressed in existing physical systems theory.

### 4.1. Ecological Interaction and Trophic Dynamics

Analyzing pufferfish role in food webs involves examining species interactions and their interrelated aspects due to the non-linear and time-variable nature of the species interaction note that variability (i.e., strength) in interaction occurs at an intrinsic level due to population dynamics; thus, if measurements only consider short measures of interaction, then they will mostly capture direct coupling and, and conversely, longer-term data would be based on indirect interactions or feedback. (Aguilar et al., 2025)

It provides some evidence to suggest that apex predators such as leopard seals utilize sophisticated system perceptions of population condition and foraging patterns to assess the impact of apex predators on ecosystem function and health (Salazar et al., 2024). However, though pufferfish are meso-predators, the challenges of monitoring their movements and population dynamics is still quite similar to those experienced when tracking marine mammals (Ngo et al., 2021). Further, the current literature on interactions does not typically consider the physical transport mechanism(s) that explain the spatial distribution of the interacting species, which is one limitation this paper will attempt to address.

### Conclusion

Pufferfish (Family Tetraodontidae) are naturally important fishes spread across marine, brackish, and freshwater environments. Their evolutionary achievement is strongly related to unique variations such as fused beak-like teeth for crushing hard-shelled prey, the capability to expand their bodies as a protection mechanism, and the addition of tetrodotoxin (TTX) for biochemical protection. These adaptations permit them to inhabit diverse trophic niches, extending from specified durophagous predators to more generalist feeders depending on habitat and life stage. Their feeding ecology shows elasticity influenced by ontogeny, season, and environmental conditions, while reserve partitioning among species decreases competition and supports co-occurrence in complex ecosystems like coral reefs, estuaries, and river systems. Behaviourally, pufferfish reveal adaptive comebacks to food obtainability and predation pressure, including hostility linked to opposition and defensive inflation when threatened. These behaviors, combined with their reproductive strategies and toxin-based defenses, highlight their combined survival strategies and significant role in aquatic food webs as both predators and prey. However, growing human impacts such as overfishing, habitat degradation, pollution, and aggressive spread in some regions pose ecological challenges. Overall, pufferfish represent an appreciated model for understanding evolutionary innovation, trophic adaptation, and predator–prey dynamics, highlighting the need for continued research and preservation efforts.

### REFERENCES

Nelson, E. J., Van Dokkum, P. G., Schreiber, N. M. F., Franx, M., Brammer, G. B., Momcheva, I. G., *et al.* (2016). Where stars form: inside-out growth and coherent star formation from HST Hα maps of 3200 galaxies across the main sequence at 0.7< z< 1.5. *The Astrophysical Journal*, *828*(1), 27.

Ulman, A., Abd Rabou, A. F. N., Al Mabruk, S., Bariche, M., Bilecenoğlu, M., Demirel, N., *et al.* (2024). Assessment of human health impacts from invasive pufferfish (attacks, poisonings and fatalities) across the Eastern Mediterranean. *Biology*, *13*(4), 208.

Marinsek, G. P., Ribeiro, C. C., Gusso-Choueri, P. K., Choueri, R. B., de Souza Abessa, D. M., & de Britto Mari, R. (2024). Multiple biomarkers in pufferfish as a proxy of environmental health in brazilian marine protected areas. *Science of The Total Environment*, *914*, 169742.

Gonçalves, G., Andriolo, U., Pinto, L., & Duarte, D. (2020). Mapping marine litter with Unmanned Aerial Systems: A showcase comparison among manual image screening and machine learning techniques. Marine pollution bulletin, 155, 111158.

Suchana, S. A., Ahmed, M. S., Islam, S. M., Rahman, M. L., Rohani, M. F., Ferdusi, T., et al. (2021). Chromium exposure causes structural aberrations of erythrocytes, gills, liver, kidney, and genetic damage in striped catfish Pangasianodon hypophthalmus. Biological Trace Element Research, 199(10), 3869-3885.

Liu, K., Sun, H., Zhao, X., Wang, C., An, C., Li, A., et al. (2024). DNA barcoding, identification, and validation of the pufferfish (Order: Tetraodontiformes) in China coastal waters. Ecology and Evolution, 14(2), e10944.

Kaplan, M., Yildirim, S., & Yildirim, D. C. (2022). Pufferfish versus lionfish: comparing risks for Turkish marine economics. Marine Economics and Management, 5(2), 173-187.

environmental health in brazilian marine protected areas. Science of The Total Environment, 914, 169742.

Otarashvili, Lasha, Subramanian, Tamilselvan, Holmberg, Jason, Levenson, J. J., & Stewart, Charles V. (2024). Multispecies Animal Re-ID Using a Large Community-Curated Dataset.

Helfman, G. S., Collette, B. B., Facey, D. E., & Bowen, B. W. (2009). The diversity of fishes: Biology, evolution, and ecology (2nd ed.). Wiley-Blackwell.

Froese, R., & Pauly, D. (2024). FishBase. World Wide Web electronic publication.
https://www.fishbase.se

Otarashvili, Lasha, Subramanian, Tamilselvan, Holmberg, Jason, Levenson, J. J., & Stewart, Charles V. (2024). Multispecies Animal Re-ID Using a Large Community-Curated Dataset.

Lasinio, Giovanna Jona, Pollice, Alessio, Marcon, Eric, & Fano, Elisa Anna (2017). Assessing the role of the spatial scale in the analysis of lagoon biodiversity. A case-study on the macrobenthic fauna of the Po River Delta. Ecological Indicators, Volume 80, September 2017, Pages 303-315, ISSN 1470-160X,

Otarashvili, Lasha, Subramanian, Tamilselvan, Holmberg, Jason, Levenson, J. J., & Stewart, Charles V. (2024). Multispecies Animal Re-ID Using a Large Community-Curated Dataset.

Otarashvili, Lasha, Subramanian, Tamilselvan, Holmberg, Jason, Levenson, J. J., & Stewart, Charles V. (2024). Multispecies Animal Re-ID Using a Large Community-Curated Dataset.

Zhang, Bowen, Boulerice, Jesse T., Kuniyil, Nikhil, Mendiratta, Charvi, Kumar, Satish, Shamon, Hila, & Manjunath, B. S. (2025). RareSpot: Spotting Small and Rare Wildlife in Aerial Imagery with Multi-Scale Consistency and Context-Aware Augmentation. https://arxiv.org/pdf/2506.19087v1
https://arxiv.org/pdf/2506.19087v1

Otarashvili, Lasha, Subramanian, Tamilselvan, Holmberg, Jason, Levenson, J. J., & Stewart, Charles V. (2024). Multispecies Animal Re-ID Using a Large Community-Curated Dataset.

Croze, Ottavio A., Martinez, Vincent A., Jakuszeit, Theresa, Dell'Arciprete, Dario, Poon, Wilson C. K., & Bees, Martin A. (2019). Helical and oscillatory microswimmer motility statistics from differential dynamic microscopy. https://doi.org/10.1088/1367-2630/ab241f https://doi.org/10.1088/1367-2630/ab241f

Lasinio, Giovanna Jona, Pollice, Alessio, Marcon, Eric, & Fano, Elisa Anna (2017). Assessing the role of the spatial scale in the analysis of lagoon biodiversity. A case-study on the macrobenthic fauna of the Po River Delta. Ecological Indicators, Volume 80, September 2017, Pages 303-315, ISSN 1470-160X,

Otarashvili, Lasha, Subramanian, Tamilselvan, Holmberg, Jason, Levenson, J. J., & Stewart, Charles V. (2024). Multispecies Animal Re-ID Using a Large Community-Curated Dataset.

Evans, K., et al. (2023). Evolutionary innovation accelerates morphological diversification in pufferfishes and their relatives. Evolution.

Renjithkumar, C. R., Roshni, K., & Ranjeet, K. (2020). Feeding ecology of the endemic freshwater puffer fish Carinotetradon travancoricus (Hora & Nair, 1941) in Western Ghats hotspot, India. International Journal of Aquatic Biology, 8(5), 300–310.

Bartolette, R., Rosa, D. C. O., Beserra, D. A., Soares, B. E., Albrecht, M. P., & Brito, M. F. G. (2018). Seasonal and ontogenetic diet patterns of the freshwater pufferfish Colomesus asellus (Müller & Troschel, 1849) in the upper-middle Tocantins River. Acta Scientiarum. Biological Sciences, 40.

Renjithkumar, C. R., Roshni, K., & Ranjeet, K. (2020). Feeding ecology of the endemic freshwater puffer fish Carinotetradon travancoricus (Hora & Nair, 1941) in Western Ghats hotspot, India. International Journal of Aquatic Biology, 8(5), 300–310.

Krumme, U., Keuthen, H., Saint-Paul, U., & Villwock, W. (1997). Contribution to the feeding ecology of the banded puffer fish Colomesus psittacus (Tetraodontidae) in north Brazilian mangrove creeks. Brazilian Journal of Biology, 67(3), 383–392.

Xie, J., et al. (2021). Effects of food quantity on aggression and monoamine levels of juvenile pufferfish (Takifugu rubripes). Behavioural Brain Research.

Ferreira, P., Kwan, G. T., Haldorson, S., Rummer, J. L., Tashiro, F., Castro, L. F. C., Tresguerres, M., & Wilson, J. M. (2022). A multi-tasking stomach: functional coexistence of acid-peptic digestion and defensive body inflation in three distantly related vertebrate lineages. Biology Letters, 18(2).

Evans, K., et al. (2023). Evolutionary innovation accelerates morphological diversification in pufferfishes and their relatives. Evolution.

McGee, G. E., & Clark, T. D. (2014). All puffed out: do pufferfish hold their breath while inflated?. Biology letters, 10(12), 20140823. https://doi.org/10.1098/rsbl.2014.0823

McGee, G. E., & Clark, T. D. (2014). All puffed out: do pufferfish hold their breath while inflated?. Biology letters, 10(12), 20140823. https://doi.org/10.1098/rsbl.2014.0823

Asano, M., Ishizaki, C., Tomonou, T., Kihara, M., Ito, M., Yasukawa, S., Shirai, K., Oyama, H., Izawa, S., Kawamura, R., Saito, K., Suo, R., Nakahigashi, R., Adachi, M., Nishikawa, T., Sugita, H., & Itoi, S. (2023). Levels of Tetrodotoxins in Spawning Pufferfish, Takifugu alboplumbeus. Marine drugs, 21(4), 207. https://doi.org/10.3390/md21040207

Smith, W. L., Stern, J. H., Girard, M. G., & Davis, M. P. (2016). Evolution of venomous cartilaginous and ray-finned fishes. Integrative and Comparative Biology, 56(5), 950–961. https://doi.org/10.1093/icb/icw070

Schoppema, M. C., Velden, B. H. M. van der, Hürriyetoğlu, A., Klijnstra, M. D., Faassen, E. J., Gerssen, A., & Fels-Klerx, H. J. van der (2025). Identifying environmental factors associated with tetrodotoxin contamination in bivalve mollusks using eXplainable AI. https://arxiv.org/pdf/2511.20395v2 https://arxiv.org/pdf/2511.20395v2

Sarkar, Trilokesh Ranjan, Das, Nilanjan, Maitra, Pralay Sankar, Some, Bijoy, Saha, Ritwik, Adhikary, Orijita, Bose, Bishal, & Sen, Jaydip (2024). Evaluating Adversarial Robustness: A Comparison Of FGSM, Carlini-Wagner Attacks, And The Role of Distillation as Defense Mechanism. https://arxiv.org/pdf/2404.04245v1 https://arxiv.org/pdf/2404.04245v1

Liu, Linbo, Park, Youngsuk, Hoang, Trong Nghia, Hasson, Hilaf, & Huan, Jun (2022). Robust Multivariate Time-Series Forecasting: Adversarial Attacks and Defense Mechanisms. https://arxiv.org/pdf/2207.09572v3 https://arxiv.org/pdf/2207.09572v3

Ge, Mengmeng, Cho, Jin-Hee, Kamhoua, Charles A., & Kim, Dong Seong (2019). Optimal Deployments of Defense Mechanisms for the Internet of Things. https://arxiv.org/pdf/1908.00324v1 https://arxiv.org/pdf/1908.00324v1

Schoppema, M. C., Velden, B. H. M. van der, Hürriyetoğlu, A., Klijnstra, M. D., Faassen, E. J., Gerssen, A., & Fels-Klerx, H. J. van der (2025). Identifying environmental factors associated with tetrodotoxin contamination in bivalve mollusks using eXplainable AI. https://arxiv.org/pdf/2511.20395v2 https://arxiv.org/pdf/2511.20395v2

Sarkar, Trilokesh Ranjan, Das, Nilanjan, Maitra, Pralay Sankar, Some, Bijoy, Saha, Ritwik, Adhikary, Orijita, Bose, Bishal, & Sen, Jaydip (2024). Evaluating Adversarial Robustness: A Comparison Of FGSM, Carlini-Wagner Attacks, And The Role of Distillation as Defense Mechanism. https://arxiv.org/pdf/2404.04245v1 https://arxiv.org/pdf/2404.04245v1

Aguilar, Javier, Suweis, Samir, Maritan, Amos, & Azaele, Sandro (2025). Unraveling the temporal dependence of ecological interaction measures. https://arxiv.org/pdf/2508.19197v1 https://arxiv.org/pdf/2508.19197v1

Selvam, A. M. (2007). Universal spectrum for DNA base CG frequency distribution in Takifugu rubripes (Puffer fish) genome. https://arxiv.org/pdf/0704.2114v2 https://arxiv.org/pdf/0704.2114v2

Selvam, A. M. (2008). Fractal Fluctuations and Statistical Normal Distribution. https://arxiv.org/pdf/0805.3426v1 https://arxiv.org/pdf/0805.3426v1

Salazar, Jorge Yero, Rivas, Pablo, Borras-Chavez, Renato, & Kienle, Sarah (2024). Image-Based Leopard Seal Recognition: Approaches and Challenges in Current Automated Systems. https://arxiv.org/pdf/2408.07269v1 https://arxiv.org/pdf/2408.07269v1

Ngô, Manh Cuong, Selvan, Raghavendra, Tervo, Outi, Heide-Jørgensen, Mads Peter, & Ditlevsen, Susanne (2021). Detection of foraging behavior from accelerometer data using U-Net type convolutional networks. https://arxiv.org/pdf/2101.01992v1 https://arxiv.org/pdf/2101.01992v1

Food and Agriculture Organization of the United Nations (FAO). (2023). *Species fact sheets: Family Tetraodontidae*. FAO Fisheries Division. https://www.fao.org

Nelson, J. S., Grande, T. C., & Wilson, M. V. H. (2016). *Fishes of the world* (5th ed.). John Wiley & Sons.

Ferreira, C. E. L., Gonçalves, J. E. A., & Coutinho, R. (2010). Feeding ecology of the checkered puffer *Sphoeroides testudineus* in Brazilian coastal waters. *Journal of Fish Biology, 76*(1), 190–204.

Noguchi, T., & Arakawa, O. (2008). Tetrodotoxin—Distribution and accumulation in aquatic organisms, and cases of human intoxication. *Marine Drugs, 6*(2), 220–242. https://doi.org/10.3390/md20080011

Yang, J., & Chen, S. (2003). Biology and artificial propagation of river pufferfish (*Takifugu obscurus*). *Journal of Applied Ichthyology, 19*(4), 233–237. https://doi.org/10.1046/j.1439-0426.2003.00470.x

Barletta, M., & Blaber, S. J. M. (2007). Comparison of fish assemblages and guilds in tropical mangrove estuaries of Brazil. *Estuarine, Coastal and Shelf Science, 71*(3–4), 513–524. https://doi.org/10.1016/j.ecss.2006.08.003

IEER
Institute for Excellence in Education & Research