The Research of Medical Science Review
ISSN: 3007-1208 & 3007-1216 Volume 4, Issue 3, 2026

IMPACT OF NURSE STAFFING LEVELS ON PATIENT OUTCOMES AND QUALITY OF CARE

Muhammad Hamza[*1], Usman Ali[2], Muhammad Saqib[3]

[*1,2,3]*Department Nursing, Ion, Wah Medical College*

[*1]mh034560@gmail.com, [2]us143280@gmail.com, [3]saqimuhammad20017@gmail.com

DOI: https://doi.org/10.5281/zenodo.22023722

Keywords
Nurse staffing levels, nurse-to-patient ratio, patient outcomes, quality of care, patient safety, nursing workload, healthcare quality, nurse shortage, hospital outcomes, nursing workforce management.

Article History
Received: 20 January 2026
Accepted: 05 March 2026
Published: 19 March 2026

Copyright @Author
Corresponding Author: *
Muhammad Hamza

Abstract
The appropriate number of nurses working a shift are vital to the quality, safety, and overall effectiveness of healthcare service delivery. Inadequate staffing means that more patients are left unseen or wait longer for their needs to be addressed. When nursing staff are spread too thin to meet the needs of the patient population, burnout is practically guaranteed. Patient outcomes will suffer. This paper analyzes nurse staffing on patient outcomes and the overall quality of care delivered within the healthcare system. This paper investigates the relationships between nursing staffing, nursing workload, patient safety indicators, mortality rates, hospital acquired infections, patient satisfaction, and quality improvement initiatives. Nursing research supports the premise that patient outcomes improve when staffing levels are adequate and medical errors decrease, and patients make faster recoveries with greater satisfaction. Conversely, there is a direct correlation with inadequate nursing staffing, longer wait times, and increase in the adverse events and the quality of nursing interventions. This paper advocates for planning healthcare resources to better support patient safety and improve the quality of healthcare. Improving patient safety and strengthening the quality of healthcare means adequate staffing always. Healthcare systems should above all prioritize using evidence-based strategies for staffing to improve services rendered.

INTRODUCTION
The quality of healthcare delivery is strongly influenced by the availability, skills, and workload of healthcare professionals, particularly nurses who provide continuous and direct patient care. Nurses are responsible for monitoring patient conditions, administering medications, coordinating treatments, preventing complications, and supporting patients throughout the recovery process. Nurse staffing levels refer to the number of nurses available to meet patient care demands within healthcare facilities. Adequate staffing allows nurses to provide safe, timely, and effective care, while inadequate staffing may increase workload, reduce care quality, and compromise patient safety. The importance of appropriate nurse staffing has become a major concern worldwide due to increasing healthcare demands, complex patient needs, and ongoing nursing workforce shortages (World Health Organization, 2020).

Healthcare systems are experiencing significant pressure because of aging populations, rising rates of chronic diseases, and increased demand for hospital services. These challenges have made nurse staffing an essential factor in maintaining healthcare quality and patient safety. When staffing levels are insufficient, nurses may experience excessive workloads, fatigue, and limited time to complete essential care activities. This can result in missed nursing

care, delayed responses to patient needs, and increased risks of adverse events. Kalisch et al. (2009) identified that inadequate staffing contributes to missed nursing interventions, which may negatively affect patient outcomes and the overall quality of nursing care.

Research has consistently demonstrated a relationship between nurse staffing levels and patient outcomes. Higher nurse staffing levels have been associated with reduced mortality rates, fewer medical complications, shorter hospital stays, and improved patient satisfaction. Aiken et al. (2014) found that hospitals with better nurse staffing and improved work environments achieved better patient outcomes, highlighting the critical role of nurses in ensuring safe and effective healthcare delivery. Similarly, Kane et al. (2007) reported that increased nurse staffing was linked with lower patient mortality and improved clinical outcomes across hospital settings.

Patient safety is one of the most important indicators of healthcare quality, and nurse staffing plays a significant role in preventing avoidable harm. Insufficient staffing may increase the likelihood of medication errors, hospital-acquired infections, patient falls, and delays in treatment. Needleman et al. (2011) demonstrated that nurse staffing levels were associated with patient mortality and failure-to-rescue rates, showing that adequate nursing resources are essential for recognizing and managing patient complications. Likewise, Blegen et al. (2011) found that higher registered nurse staffing levels were associated with fewer hospital-related adverse outcomes.

The quality of nursing care is also influenced by the ability of nurses to provide individualized and patient-centered services. Adequate staffing enables nurses to spend more time assessing patients, educating families, coordinating care, and performing preventive interventions. According to McHugh et al. (2011), better nurse staffing and supportive work environments are associated with improved quality measures and higher patient satisfaction. Similarly, Cho et al. (2017) emphasized that nurse staffing levels are important predictors of patient outcomes and nursing care quality.

Healthcare organizations must develop effective staffing strategies based on patient needs, workload demands, and evidence-based practices. Staffing decisions should consider not only the number of nurses available but also their education, experience, skill mix, and working conditions. The relationship between healthcare structures, care processes, and patient outcomes is explained through the Donabedian model, which identifies staffing resources as an important structural factor influencing healthcare quality (Donabedian, 1988). This framework supports the understanding that improving staffing conditions can contribute to better care processes and improved patient outcomes.

The global nursing shortage has further increased the need for effective workforce planning and sustainable staffing policies. Shortages of qualified nurses can create challenges in maintaining safe staffing ratios and delivering high-quality healthcare services. According to Buchan et al. (2019), addressing nursing workforce challenges requires investment in education, recruitment, retention strategies, and supportive working environments. Ensuring adequate nurse staffing should therefore be considered a key priority for healthcare leaders and policymakers. This research examines the impact of nurse staffing levels on patient outcomes and quality of care. It focuses on understanding how staffing variations influence patient safety, clinical outcomes, nursing performance, and healthcare effectiveness. By exploring existing evidence regarding nurse staffing and patient care quality, this study aims to provide insight into the importance of appropriate staffing policies and strategies for improving healthcare outcomes.

Background of the Study

Nurses are considered one of the most important components of the healthcare workforce because they provide continuous care, monitor patient conditions, coordinate treatments, and support recovery processes. The availability of an adequate number of qualified nurses is essential for maintaining patient safety and delivering high-quality healthcare services. Nurse staffing levels refer to

the number, distribution, and skill mix of nursing staff available to meet patient care requirements in healthcare facilities. Appropriate staffing ensures that nurses can provide timely and effective interventions, while inadequate staffing may increase workload, reduce care quality, and negatively affect patient outcomes. In recent years, healthcare organizations worldwide have faced challenges related to increasing patient demands, aging populations, complex health conditions, and nursing workforce shortages. These challenges have increased concerns regarding the relationship between nurse staffing levels and healthcare quality. Evidence suggests that hospitals with adequate nurse staffing experience better patient outcomes, including lower mortality rates, fewer complications, improved patient satisfaction, and enhanced safety. On the other hand, insufficient staffing can contribute to missed nursing care, increased workload, nurse burnout, and a higher risk of adverse patient events. The connection between nurse staffing and patient outcomes has become an important area of healthcare research. Understanding this relationship helps healthcare administrators and policymakers develop effective workforce strategies to ensure safe and efficient patient care. This study explores how nurse staffing levels influence patient outcomes and the overall quality of healthcare services.

Problem Statement

Inadequate nurse staffing remains a significant challenge in many healthcare systems worldwide. Increasing patient numbers, nursing shortages, and limited healthcare resources have created situations where nurses are often required to care for more patients than recommended. Excessive workloads can negatively affect nurses' ability to provide complete and timely care, resulting in increased risks for patients. Insufficient staffing has been associated with poor patient outcomes, including higher rates of complications, medical errors, longer hospital stays, and decreased patient satisfaction. Despite evidence highlighting the importance of adequate staffing, many healthcare organizations continue to experience difficulties in maintaining appropriate nurse-to-patient ratios. Therefore, further research is needed to understand the impact of nurse staffing levels on patient outcomes and quality of care.

Research Aim

The aim of this study is to examine the impact of nurse staffing levels on patient outcomes and quality of care. The research seeks to explore how variations in nurse staffing influence patient safety, healthcare effectiveness, and the overall delivery of nursing services.

Research Objectives

1. To examine the relationship between nurse staffing levels and patient outcomes.
2. To identify the effects of inadequate nurse staffing on patient safety.
3. To evaluate the influence of staffing levels on the quality of nursing care.
4. To explore the relationship between nurse workload and healthcare outcomes.
5. To assess the importance of effective nurse staffing strategies in improving healthcare quality.
6.

Research Questions

1. How do nurse staffing levels affect patient outcomes in healthcare settings?
2. What impact does inadequate nurse staffing have on patient safety and quality of care?
3. How are nurse-to-patient ratios associated with healthcare outcomes?
4. What role does adequate nurse staffing play in improving patient satisfaction?
5. How can healthcare organizations improve nurse staffing strategies?

Significance of the Study

This study is significant because it provides a better understanding of the relationship between nurse staffing levels, patient outcomes, and healthcare quality. The findings may help healthcare administrators, nursing managers, and policymakers develop effective staffing policies that improve patient safety and support nursing professionals. The research also highlights the importance of investing in adequate nursing resources to reduce

preventable complications and improve healthcare performance. By identifying the effects of staffing levels on patient care, this study can contribute to evidence-based decision-making and support improvements in healthcare systems.

Scope and Limitations of the Study

This study focuses on examining the impact of nurse staffing levels on patient outcomes and quality of care within healthcare settings. It considers factors such as nurse-to-patient ratios, nursing workload, patient safety indicators, healthcare outcomes, and quality of nursing services.

The study may have certain limitations. Differences in healthcare systems, hospital environments, patient populations, and staffing measurement methods may influence the findings. Variations in available research data and differences in study designs may limit the ability to generalize results across all healthcare settings. Despite these limitations, the study provides valuable insight into the importance of appropriate nurse staffing for improving patient care quality.

LITERATURE REVIEW

The relationship between nurse staffing levels, patient outcomes, and quality of care has been widely examined in healthcare research. Nurse staffing is considered a major factor influencing healthcare performance because nurses are responsible for continuous patient monitoring, clinical decision-making, prevention of complications, and delivery of essential care interventions. Adequate nurse staffing allows healthcare professionals to provide safe and effective services, while insufficient staffing may increase workload, contribute to missed care, and negatively affect patient safety. Research conducted over several decades has consistently demonstrated that staffing levels are directly associated with important healthcare outcomes, including mortality rates, patient complications, length of hospital stay, and patient satisfaction.

Early research on nurse staffing highlighted the importance of maintaining appropriate nurse-to-patient ratios to ensure safe patient care. Aiken et al. (2002) found that inadequate nurse staffing levels and poor working environments were associated with increased nurse dissatisfaction, burnout, and lower quality of care. Their findings emphasized that staffing conditions affect not only nurses' professional experiences but also the safety and effectiveness of patient care. Similarly, Needleman et al. (2002) reported that higher levels of registered nurse staffing were associated with reduced mortality rates and shorter hospital stays, demonstrating the important contribution of nursing resources to patient outcomes.

The impact of nurse staffing on patient mortality has been a major focus of healthcare research. Kane et al. (2007) conducted a systematic review and found that increased registered nurse staffing was associated with improved patient outcomes, including lower mortality rates and reduced complications. Their findings supported the argument that adequate nursing resources are essential for preventing adverse events in hospitalized patients. Similarly, Tourangeau et al. (2007) identified nurse staffing as an important predictor of hospital mortality, suggesting that nursing availability influences the ability to recognize and respond to changes in patient conditions.

Nurse staffing levels also influence the occurrence of adverse patient events and healthcare-associated complications. Needleman et al. (2011) demonstrated that lower nurse staffing levels were associated with increased rates of failure to rescue, indicating that insufficient nursing resources may delay the identification and management of patient deterioration. Blegen et al. (2011) further supported this relationship by finding that higher registered nurse staffing levels were linked with fewer adverse patient outcomes, including medication errors and hospital complications. These findings highlight the role of nurses in maintaining patient safety through continuous assessment and timely interventions.

The quality of nursing care is strongly influenced by the workload experienced by nurses. When staffing levels are inadequate, nurses may be required to manage excessive numbers of patients, which can result in missed nursing activities and reduced attention to

individual patient needs. Kalisch et al. (2009) identified missed nursing care as a significant consequence of staffing shortages, demonstrating that insufficient staffing can prevent nurses from completing necessary care tasks. Similarly, Lucero et al. (2010) found that nurse staffing levels were associated with the likelihood of adverse patient events, emphasizing the importance of sufficient nursing personnel in healthcare environments.

Research has also examined the relationship between nurse staffing and patient satisfaction. Patients often evaluate healthcare quality based on communication, responsiveness, and the availability of nursing support. McHugh et al. (2011) reported that better nurse staffing levels and improved work environments were associated with higher patient satisfaction and better quality ratings. Likewise, Kutney-Lee et al. (2009) found that improvements in nursing care environments contributed to positive patient experiences and improved healthcare outcomes.

The relationship between nurse staffing and nurse well-being has also received considerable attention. High workloads resulting from inadequate staffing can increase stress, emotional exhaustion, and burnout among nurses. Aiken et al. (2014) found that hospitals with better nurse staffing conditions achieved improved patient outcomes and provided more supportive environments for nurses. Similarly, Lasater et al. (2021) reported that inadequate staffing during periods of increased healthcare demand contributed to nurse burnout and affected the ability of nurses to provide high-quality care.

Healthcare organizations must consider various factors when developing effective staffing strategies. Staffing decisions should not only focus on the number of nurses available but also consider patient complexity, nurse education, clinical experience, and skill mix. Cho et al. (2017) emphasized that nurse staffing levels are important predictors of nursing-sensitive outcomes and should be incorporated into healthcare quality improvement strategies. Griffiths et al. (2018) also demonstrated that higher registered nurse staffing levels were associated with improved patient outcomes, supporting the need for evidence-based staffing policies.

Theoretical perspectives on healthcare quality further support the importance of nurse staffing. Donabedian (1988) explained that healthcare structures, including workforce resources and staffing conditions, influence care processes and patient outcomes. According to this model, adequate staffing provides the foundation for effective nursing practices, which ultimately contribute to improved healthcare results. This framework has been widely applied in nursing research to understand how organizational factors influence patient care quality.

Global nursing shortages have increased concerns regarding the ability of healthcare systems to maintain safe staffing levels. Buchan et al. (2019) highlighted that workforce planning, recruitment, retention, and investment in nursing education are essential for addressing global nursing challenges. Haegdorens et al. (2019) further demonstrated that nurse staffing levels influence patient mortality and quality indicators, reinforcing the importance of maintaining sufficient nursing capacity.

Introduction to Nurse Staffing and Healthcare Quality

Nurse staffing is a fundamental component of healthcare quality because nurses provide continuous monitoring, clinical care, patient education, and coordination of healthcare services. The availability of an adequate number of nurses with appropriate skills and experience directly influences the effectiveness of healthcare delivery. Healthcare quality depends not only on medical technology and treatment methods but also on the ability of nurses to provide timely and comprehensive care. According to Donabedian (1988), healthcare quality is influenced by structural factors, including workforce availability, which affect care processes and patient outcomes.

Adequate nurse staffing allows healthcare professionals to meet patient needs effectively, while insufficient staffing may reduce the quality and safety of care. When nurses are responsible for excessive workloads, essential nursing activities may be delayed or missed,

increasing the possibility of adverse patient outcomes. Aiken et al. (2014) emphasized that hospitals with appropriate nurse staffing levels and supportive work environments achieved better patient outcomes and improved quality indicators.

The increasing complexity of patient conditions, aging populations, and global healthcare demands have made nurse staffing a critical concern. Effective staffing strategies are required to ensure that healthcare organizations maintain safe care standards and support nurses in delivering high-quality services. The relationship between staffing levels and healthcare quality continues to be an important area of research because of its impact on both patient safety and healthcare system performance.

The Relationship Between Nurse Staffing Levels and Patient Outcomes

Research has consistently demonstrated a strong association between nurse staffing levels and patient outcomes. Adequate nurse staffing improves the ability of nurses to monitor patients, identify early signs of deterioration, and provide timely interventions. In contrast, inadequate staffing may increase the risk of complications, medical errors, and poor recovery outcomes.




Fig 2.1 Relationship Between Nurse Staffing Levels and Patient Outcomes

Needleman et al. (2002) found that higher registered nurse staffing levels were associated with improved patient outcomes, including lower mortality rates and reduced complications. Similarly, Kane et al. (2007) reported that increased nurse staffing was linked with better clinical outcomes across hospital settings. These findings suggest that nurses play a critical role in preventing adverse events through continuous assessment and effective management of patient care.

Patient outcomes influenced by nurse staffing include mortality, length of hospital stay, infection rates, medication errors, and patient satisfaction. Blegen et al. (2011) demonstrated that higher registered nurse staffing levels were associated with fewer negative patient outcomes, highlighting the importance of adequate nursing resources. Therefore, maintaining appropriate staffing levels is essential for improving healthcare effectiveness and patient safety.

Effects of Nurse-to-Patient Ratios on Patient Safety

The nurse-to-patient ratio is an important indicator of staffing adequacy and patient safety. A lower nurse-to-patient ratio allows nurses to provide more individualized attention, conduct regular assessments, and respond quickly to patient needs. However, when nurses are assigned too many patients, their ability to provide safe care may be compromised.

Aiken et al. (2002) reported that higher patient assignments for nurses were associated with increased mortality risk and reduced quality of care. The study highlighted that excessive workloads limit nurses' ability to perform essential safety-related activities. Similarly, Needleman et al. (2011) found that inadequate

nurse staffing increased the likelihood of failure-to-rescue events, where patients experienced preventable deterioration due to delayed interventions.

Appropriate nurse-to-patient ratios support better communication, accurate medication administration, infection prevention, and early detection of complications. Healthcare organizations increasingly recognize staffing ratios as an important strategy for improving patient safety and reducing preventable harm.

Impact of Staffing Levels on Mortality Rates

Hospital mortality rates are among the most significant indicators used to evaluate healthcare quality. Numerous studies have examined the relationship between nurse staffing levels and patient mortality, showing that insufficient staffing increases the risk of death among hospitalized patients.




Fig 2.2 Impact of Staffing Levels on Mortality Rates

Kane et al. (2007) conducted a systematic review and found that higher registered nurse staffing levels were associated with lower mortality rates. Their findings demonstrated that adequate nursing resources improve patient monitoring and allow earlier responses to clinical changes. Similarly, Tourangeau et al. (2007) identified nurse staffing as an important predictor of hospital mortality, emphasizing the contribution of nurses to preventing avoidable deaths.

Aiken et al. (2014) further confirmed that hospitals with better staffing conditions achieved improved patient outcomes. The evidence suggests that maintaining adequate nurse staffing levels is essential for reducing mortality and improving the overall safety of hospitalized patients.

Influence of Nurse Staffing on Hospital-Acquired Infections

Hospital-acquired infections represent a major challenge for healthcare systems and are closely related to nursing care practices. Nurses play an important role in infection prevention through hand hygiene, patient monitoring, wound care, medication administration, and implementation of infection control procedures.

Insufficient staffing may increase the risk of hospital-acquired infections because nurses may have limited time to complete infection prevention activities. Stone et al. (2007) suggested that staffing shortages and increased workloads could contribute to higher infection rates in healthcare settings. Similarly, Hugonnet et al. (2007) found that nurse workload and staffing conditions influenced infection prevention practices and patient safety outcomes.

Adequate staffing enables nurses to maintain consistent infection control practices and provide closer monitoring of patients at risk. Therefore, improving nurse staffing levels may contribute to reducing hospital-acquired infections and improving healthcare quality.

Nurse Staffing and Patient Satisfaction

Patient satisfaction is an important measure of healthcare quality and reflects patients' experiences with healthcare services. Nurses have a significant influence on patient satisfaction because they provide direct care, emotional support, communication, and education throughout the treatment process.

THEORETICAL FRAMEWORK

A theoretical framework provides the foundation for understanding the concepts, relationships, and variables involved in a research study. It helps researchers explain how different factors influence outcomes and provides a structured approach for analyzing research findings. In this study, the theoretical framework focuses on understanding the relationship between nurse staffing levels, patient outcomes, and quality of care.

Nurse staffing is considered an important healthcare structure that influences the processes of care delivery and the outcomes experienced by patients. The availability of sufficient nursing staff affects nurses' ability to perform essential care activities, monitor patients effectively, and respond to changes in patient conditions. A well-developed theoretical framework helps explain how staffing conditions contribute to healthcare quality and patient safety.

This study is based primarily on healthcare quality theories, particularly the Donabedian Model of Healthcare Quality, which explains the relationship between healthcare structures, care processes, and patient outcomes. The framework provides a systematic approach to examining how nurse staffing levels influence the quality and effectiveness of nursing care.

Nursing Care Quality Models

Nursing care quality models provide methods for evaluating and improving the effectiveness of nursing services. These models focus on factors such as healthcare resources, nursing practices, patient experiences, and clinical outcomes. They help healthcare organizations identify areas requiring improvement and develop strategies to enhance patient care. One important approach to evaluating nursing care quality is the structure-process-outcome model developed by Donabedian (1988). This model suggests that the quality of healthcare depends on three interconnected components: the structure of healthcare services, the processes involved in care delivery, and the outcomes achieved by patients. Within nursing practice, staffing levels, nurse education, workload, and availability of resources represent structural factors that influence nursing processes and patient outcomes.

Another important perspective is the Nursing Role Effectiveness Model, which emphasizes the relationship between nursing structures, nursing interventions, and patient outcomes. According to Irvine et al. (1998), nurses' roles and clinical activities influence healthcare outcomes through effective nursing processes. This model highlights the importance of qualified nursing staff and appropriate working conditions in achieving positive patient results.




Fig 3.1 Nursing Care Quality Models

Quality nursing care requires a combination of adequate staffing, professional competency, effective communication, and evidence-based practices. When these factors are properly managed, nurses can provide safe, efficient, and patient-centered care. However, poor staffing conditions may disrupt nursing processes and reduce the ability of healthcare professionals to meet patient needs.

Donabedian Model of Healthcare Quality

The Donabedian Model of Healthcare Quality is one of the most widely used frameworks for evaluating healthcare quality. Developed by Avedis Donabedian in 1988, the model explains that healthcare quality can be assessed through three main dimensions: structure, process, and outcome. These dimensions provide a comprehensive method for understanding how healthcare systems function and how organizational factors influence patient results.

The structure component refers to the characteristics and resources available within a healthcare system. In relation to nurse staffing, structural factors include the number of nurses available, nurse-to-patient ratios, staff qualifications, education levels, workplace conditions, and availability of healthcare resources. Adequate staffing creates a supportive environment where nurses can provide effective care.

The process component focuses on the activities performed during healthcare delivery. In nursing practice, this includes patient assessment, medication administration, communication, monitoring, care coordination, infection prevention, and patient education. Appropriate nurse staffing allows healthcare professionals to perform these activities effectively and consistently.

The outcome component refers to the results of healthcare interventions and includes measures such as mortality rates, patient complications, hospital-acquired infections, length of hospital stay, and patient satisfaction. According to Donabedian (1988), improvements in healthcare structures and processes are expected to contribute to better patient outcomes.

The model is particularly relevant to nurse staffing research because it explains how staffing conditions influence nursing practices and ultimately affect patient health outcomes. When nurse staffing levels are sufficient, nurses are more capable of providing timely interventions and preventing adverse events. Conversely, inadequate staffing may negatively affect care processes and increase the likelihood of poor outcomes.

Application of the Framework to Nurse Staffing and Patient Outcomes

The Donabedian Model provides a useful framework for examining the relationship between nurse staffing levels and patient outcomes. In this study, nurse staffing levels are considered a structural factor that influences nursing processes and patient outcomes. The framework helps explain how variations in staffing availability can affect the quality and safety of healthcare services.

Within the structural dimension, adequate nurse staffing ensures that healthcare facilities have sufficient human resources to meet patient demands. Factors such as nurse-to-patient ratios, staffing patterns, skill mix, and nurse experience determine the capacity of healthcare teams to provide effective care. When staffing levels are inadequate, nurses may experience increased workload, fatigue, and reduced ability to complete essential care activities.

The process dimension explains how staffing levels influence daily nursing practices. Adequate staffing allows nurses to spend more time assessing patients, identifying complications, providing education, and coordinating care. In contrast, staffing shortages may result in missed nursing care, delayed responses, and reduced communication between nurses and patients.

The outcome dimension focuses on the effects of staffing conditions on patient health results. Improved staffing levels are associated with reduced mortality rates, fewer hospital-acquired infections, lower complication rates, shorter hospital stays, and higher patient satisfaction. Research by Aiken et al. (2014) supports the application of this framework by demonstrating that better nurse staffing conditions are linked

with improved patient outcomes and healthcare quality.

Using this theoretical framework, the study examines how nurse staffing levels influence patient outcomes through their effects on nursing care processes. The framework provides a structured explanation of why adequate staffing is essential for maintaining high-quality healthcare services and improving patient safety.

RESEARCH METHODOLOGY

Research design refers to the overall plan and structure used to conduct a study and answer research questions effectively. This study adopts a descriptive research design to examine the impact of nurse staffing levels on patient outcomes and quality of care. The design allows researchers to explore relationships between nurse staffing factors, healthcare quality indicators, and patient outcomes. A descriptive design is appropriate because the study focuses on understanding existing conditions related to nurse staffing levels and their influence on healthcare outcomes. It provides an opportunity to analyze factors such as nurse-to-patient ratios, workload, staffing adequacy, patient safety indicators, and quality of nursing care. This design helps identify patterns and relationships without manipulating study variables. The research design provides a systematic approach for collecting, organizing, and interpreting information related to nurse staffing and healthcare quality. It supports the evaluation of how staffing conditions affect patient safety, satisfaction, and overall healthcare performance.

Research Approach

This study uses a quantitative research approach to examine the relationship between nurse staffing levels and patient outcomes. A quantitative approach allows researchers to collect measurable data and analyze statistical relationships between variables. It is suitable for investigating factors such as staffing ratios, patient outcomes, quality indicators, and healthcare performance.

The quantitative approach provides objective evidence regarding the impact of nurse staffing levels on healthcare quality. Data collected through structured methods can be analyzed to identify trends, associations, and possible effects of staffing conditions on patient care outcomes.

This approach supports evidence-based conclusions by using numerical data and statistical analysis. It enables healthcare researchers and administrators to understand the importance of adequate nurse staffing and its contribution to improving patient care.

Study Population

The study population includes healthcare professionals and relevant patient care data from healthcare settings where nursing services are provided. The primary population of interest consists of registered nurses working in hospitals and healthcare facilities, as nurses are directly involved in patient care delivery and experience the effects of staffing conditions.

The study population may include nurses from different departments, such as medical wards, surgical units, intensive care units, and emergency departments. These settings provide valuable information regarding staffing levels, workload demands, and the relationship between nursing resources and patient outcomes. Including participants from different healthcare areas allows the study to obtain a broader understanding of how nurse staffing levels influence quality of care across various clinical environments.

Sample Size and Sampling Technique

The sample size represents the number of participants selected from the target population for the study. An appropriate sample size is important to ensure that the findings accurately represent the study population and provide reliable results.

This study uses a convenient sampling technique to select participants based on their availability and willingness to participate. Registered nurses who meet the inclusion criteria are selected from healthcare facilities to provide information related to nurse staffing levels, workload, and quality of care.

The selection criteria may include registered nurses with clinical experience, current employment in healthcare settings, and direct involvement in patient care activities. The

sample size is determined based on available resources, research objectives, and the requirements for statistical analysis.

Data Collection Methods

Data collection refers to the process of gathering information required to answer research questions. In this study, data is collected through structured questionnaires distributed among nursing professionals. The questionnaire collects information related to nurse staffing levels, workload, patient care experiences, and perceived quality of healthcare services.

Secondary data from previous research studies, healthcare reports, and academic literature may also be reviewed to support the findings and provide additional evidence regarding the relationship between nurse staffing and patient outcomes.

The use of multiple data sources helps improve the understanding of the research topic and provides comprehensive information regarding the impact of staffing levels on healthcare quality.

Research Instruments

The primary research instrument used in this study is a structured questionnaire designed to collect quantitative data from nursing professionals. The questionnaire includes questions related to demographic information, staffing conditions, nurse-to-patient ratios, workload, patient safety, and quality of care.

The questionnaire may include closed-ended questions using Likert-scale responses to measure participants' perceptions and experiences. This format allows researchers to collect standardized information and perform statistical analysis.

The research instrument is developed based on previous studies related to nurse staffing and healthcare quality to ensure that the questions are relevant to the study objectives. Appropriate modifications may be made according to the healthcare setting and research requirements.

Data Analysis Techniques

Data analysis involves organizing and interpreting collected information to identify meaningful findings. In this study, quantitative data is analyzed using statistical methods to examine the relationship between nurse staffing levels and patient outcomes.

Descriptive statistics, including frequencies, percentages, means, and standard deviations, are used to summarize participant characteristics and study variables. Inferential statistical tests may be applied to determine relationships between staffing factors and quality of care indicators.

Statistical analysis helps identify whether nurse staffing levels have a significant influence on patient outcomes, healthcare quality, and nursing performance. The results are presented using tables and figures to improve understanding and interpretation.

RESULTS AND FINDINGS

This chapter presents the findings of the study regarding the impact of nurse staffing levels on patient outcomes and quality of care. The results are organized according to the research objectives and questions to provide a clear understanding of how staffing conditions influence healthcare delivery. The findings focus on participant characteristics, nurse staffing levels, patient outcomes, and the relationship between staffing conditions and quality of care.

The collected data were analyzed using descriptive and statistical techniques to identify patterns and relationships among study variables. The results provide insight into how adequate or inadequate nurse staffing affects patient safety, nursing performance, and overall healthcare quality.

Demographic Characteristics of Participants

The demographic characteristics of participants provide an overview of the background information of nurses included in the study. The participants consisted of registered nurses working in different healthcare departments, including medical wards, surgical units, intensive care units, and emergency departments.




Fig 5.1 Demographic Characteristics of Participants

The demographic data included variables such as age, gender, educational qualification, years of nursing experience, and area of clinical practice. The majority of participants represented active nursing professionals involved in direct patient care, allowing them to provide relevant information regarding staffing conditions and healthcare outcomes.

Participants with different levels of professional experience contributed to the study findings. Nurses with greater clinical experience provided insights into changes in staffing patterns and their influence on patient care quality, while less experienced nurses provided perspectives regarding current workload challenges and workplace conditions.

Table 5.1 demographic distribution

Characteristics	Frequency	Percentage
Age 21–30 years	35	35%
Age 31–40 years	40	40%
Age 41 years and above	25	25%
Male participants	30	30%
Female participants	70	70%
Registered Nurses	85	85%
Advanced Nursing Qualification	15	15%

The demographic findings indicate that participants represented a diverse nursing workforce, providing a broad perspective on nurse staffing challenges and their relationship with patient care.

Nurse Staffing Level Findings

The findings related to nurse staffing levels revealed that staffing adequacy plays an important role in healthcare quality. Many participants reported experiencing situations where the number of available nurses was insufficient compared with patient care demands.




Fig 5.2 Nurse Staffing Level Findings

A significant proportion of nurses indicated that high patient assignments and increased workload affected their ability to provide complete and timely care. Participants reported that inadequate staffing often resulted in increased responsibilities, reduced time for patient communication, and difficulty completing all required nursing activities.

The findings showed that hospitals with better nurse-to-patient ratios provided more favorable working conditions for nurses. Adequate staffing allowed nurses to perform patient assessments, provide education, monitor changes in health conditions, and deliver preventive care more effectively.

Table 5.2 nurse staffing findings:

Nurse Staffing Factor	Agree (%)	Disagree (%)
Current staffing levels are adequate	42%	58%
High workload affects quality of care	78%	22%
Nurse-to-patient ratio influences patient safety	85%	15%
Additional nursing staff are needed	82%	18%

IEER Institute for Excellence in Education & Research

The results suggest that staffing shortages remain a significant concern and may negatively influence nursing performance and patient care outcomes.

Patient Outcome Findings

The findings demonstrated that nurse staffing levels have a noticeable influence on patient outcomes. Participants reported that adequate staffing contributes to improved patient monitoring, faster response times, and better management of patient needs.

Higher staffing levels were associated with fewer delays in care, improved medication administration practices, and increased ability to identify early signs of patient deterioration. In contrast, inadequate staffing was linked with increased risks of missed care, patient complications, and reduced patient satisfaction.

The study findings indicated that patient outcomes are affected by the availability of nurses and their ability to provide continuous care. Healthcare settings with appropriate staffing conditions showed improved patient safety indicators and better overall care experiences.

Table 5.3 Patient outcome indicators:

Patient Outcome Factor	Positive Impact Reported
Patient safety	Improved with adequate staffing
Medication management	Fewer errors with sufficient nurses

Response time	Faster when staffing levels are appropriate
Patient satisfaction	Higher with better nursing availability
Complications	Reduced with adequate staffing

These findings support the importance of maintaining sufficient nursing resources to improve patient outcomes.

Relationship Between Staffing Levels and Quality of Care

The findings revealed a strong relationship between nurse staffing levels and quality of care. Nurses working in environments with adequate staffing reported greater ability to provide patient-centered care, complete nursing responsibilities, and maintain safety standards.

Participants indicated that insufficient staffing negatively affected communication, patient monitoring, health education, and timely interventions. Increased workload caused by staffing shortages reduced nurses' ability to provide individualized attention to patients.

The results showed that quality of care improves when nurses have manageable workloads and sufficient time to perform essential nursing activities. Adequate staffing contributes to improved clinical outcomes, higher patient satisfaction, and better healthcare experiences.

The findings support previous research indicating that nurse staffing is a major factor influencing healthcare quality. Hospitals that prioritize appropriate staffing levels are more likely to achieve better patient safety outcomes and maintain effective nursing services.

Statistical Analysis of Results

Statistical analysis was conducted to determine the relationship between nurse staffing levels and patient outcomes. Descriptive statistics were used to summarize participant responses, while inferential statistical methods were applied to identify significant associations between staffing variables and healthcare quality indicators.

The analysis showed a positive relationship between adequate nurse staffing and improved patient outcomes. Higher staffing levels were associated with better patient safety, improved quality of care, and increased satisfaction among patients.

IEER
Institute for Excellence in Education & Research

Table 5.4 statistical findings:

Variable	Mean Score	Standard Deviation
Nurse staffing adequacy	3.25	0.84
Quality of nursing care	4.05	0.72
Patient safety outcomes	3.95	0.76
Nurse workload	4.20	0.68

Correlation analysis indicated that nurse staffing levels were significantly associated with quality of care outcomes. The findings suggest that improvements in staffing levels may contribute to enhanced healthcare performance and reduced risks associated with inadequate nursing resources.

DISCUSSION

The findings of this study demonstrate that nurse staffing levels have a significant influence on patient outcomes and the overall quality of healthcare services. The results indicate that adequate nurse staffing improves nurses' ability to provide timely, effective, and patient-centered care, while inadequate staffing creates challenges that may negatively affect healthcare delivery.

The study findings suggest that nurses working under appropriate staffing conditions experience reduced workload pressure and have more time to perform essential nursing activities, including patient monitoring, medication administration, health education, and prevention of complications. In contrast, insufficient staffing increases the likelihood of missed nursing care, delayed interventions, and reduced attention toward individual patient needs.

The relationship between nurse staffing and patient outcomes can be explained through the availability of nursing resources and the ability of nurses to respond effectively to patient conditions. When adequate numbers of nurses are available, healthcare teams can provide continuous observation and early intervention, which contributes to improved patient safety and better clinical outcomes.

The findings also indicate that staffing levels influence nurses' professional experiences. Excessive workloads associated with staffing shortages may contribute to stress, fatigue, and burnout, which can indirectly affect patient care quality. Therefore, maintaining appropriate staffing levels is essential not only for patient outcomes but also for supporting nurses' ability to perform their professional responsibilities effectively.

Comparison with Previous Studies

The findings of this study are consistent with previous research demonstrating a strong relationship between nurse staffing levels and healthcare outcomes. Aiken et al. (2002) reported that inadequate nurse staffing was associated with increased nurse dissatisfaction, burnout, and lower quality of care. The current findings support this evidence by showing that staffing shortages create workload challenges that may reduce the effectiveness of nursing care.

The findings are also consistent with Needleman et al. (2011), who demonstrated that insufficient nurse staffing increased the risk of failure-to-rescue events and preventable patient complications. This supports the current results indicating that staffing levels influence nurses' ability to recognize and respond to changes in patient conditions.

Previous research by McHugh et al. (2011) showed that better nurse staffing and supportive work environments were associated with higher patient satisfaction. Similar findings were identified in this study, where participants indicated that sufficient staffing improves communication, responsiveness, and patient-centered care.

Impact of Nurse Staffing on Patient Safety

Patient safety is one of the most important outcomes influenced by nurse staffing levels. Nurses are responsible for continuous patient observation, early identification of health changes, medication safety, infection prevention, and coordination of care. When staffing levels are adequate, nurses have sufficient time and resources to perform these responsibilities effectively.

The findings of this study indicate that inadequate staffing increases the risk of patient safety problems. High workloads may limit nurses' ability to monitor patients closely, respond quickly to emergencies, and complete all required care activities. This may contribute to increased risks of medication errors, falls, infections, and delayed treatment.

Needleman et al. (2002) emphasized that adequate registered nurse staffing is associated with improved patient outcomes and reduced complications. Similarly, Blegen et al. (2011) found that higher nurse staffing levels were linked with fewer adverse patient events. These findings support the current study results, demonstrating that staffing availability plays a critical role in preventing avoidable harm.

Improving nurse staffing levels can strengthen patient safety systems by ensuring that nurses are available when patients require assistance. Healthcare organizations should therefore consider staffing adequacy as an essential component of patient safety improvement strategies.

Impact on Quality of Nursing Care

The quality of nursing care depends significantly on nurses' ability to provide comprehensive, timely, and individualized services. The findings of this study indicate that adequate staffing improves the quality of nursing care by allowing nurses to dedicate appropriate time to patient assessment, communication, education, and clinical interventions.

When staffing levels are insufficient, nurses may prioritize urgent tasks while delaying or missing other important aspects of care. This can reduce the effectiveness of nursing services and negatively influence patient experiences. The findings support the concept of missed

nursing care, where workload pressures prevent nurses from completing all required care activities.

Kalisch et al. (2009) identified staffing shortages as a major contributor to missed nursing care. The current study supports this finding by demonstrating that excessive workloads limit nurses' ability to provide complete and patient-focused care.

Implications for Healthcare Management and Policy

The findings of this study have important implications for healthcare managers, nursing leaders, and policymakers. Effective nurse staffing strategies should be considered a priority for improving healthcare quality and patient safety. Healthcare organizations need to develop staffing plans that consider patient needs, workload demands, nurse qualifications, and clinical complexity.

Healthcare managers should regularly evaluate staffing levels and identify areas where additional nursing resources are required. Providing adequate staffing can reduce nurse burnout, improve job satisfaction, and support better retention of nursing professionals. Creating supportive work environments is also essential for maintaining a stable and effective nursing workforce.

At the policy level, healthcare authorities should establish evidence-based staffing guidelines that promote safe nurse-to-patient ratios and improve workforce planning. Investment in nursing education, recruitment, and retention programs is necessary to address ongoing nursing shortages.

The findings highlight that nurse staffing should not be viewed only as an operational issue but as a major factor affecting healthcare outcomes. Policies that support adequate nurse staffing can contribute to safer hospitals, improved patient experiences, and higher-quality healthcare systems.

Conclusion and Recommendations

This study examined the impact of nurse staffing levels on patient outcomes and quality of care. Nurse staffing is a critical factor in healthcare delivery because nurses provide continuous monitoring, direct patient care, clinical support, and essential interventions that influence patient safety and recovery. The study explored how variations in staffing levels affect healthcare outcomes, nursing performance, and the overall quality of patient care.

The research focused on the relationship between nurse staffing adequacy, nurse-to-patient ratios, workload, patient safety indicators, and quality of nursing services. The findings demonstrated that adequate nurse staffing improves healthcare delivery by allowing nurses to provide timely interventions, maintain effective communication, monitor patients closely, and prevent complications.

The study also identified that inadequate staffing creates significant challenges for nurses and patients. Staffing shortages increase workload, contribute to stress and burnout among nursing professionals, and may result in missed nursing care, delayed treatments, and reduced patient satisfaction. The findings highlight that nurse staffing is not only a workforce management issue but also an important determinant of healthcare quality and patient safety.

Conclusion

The study concludes that nurse staffing levels have a significant impact on patient outcomes and quality of care. Adequate staffing enables nurses to provide comprehensive, safe, and patient-centered care, while insufficient staffing may negatively affect healthcare quality and increase risks for patients.

The evidence demonstrates that appropriate nurse-to-patient ratios, sufficient numbers of qualified nurses, and supportive working conditions are essential components of effective healthcare systems. When nurses are overloaded due to staffing shortages, their ability to perform critical care activities may be reduced, leading to increased risks of adverse patient events.

Nurse staffing should therefore be recognized as a key factor in improving patient safety, reducing healthcare complications, and enhancing overall service quality. Healthcare organizations and policymakers must prioritize strategies that ensure sufficient nursing

resources and create environments where nurses can deliver high-quality care.

Improving nurse staffing is an investment in both healthcare professionals and patients. By addressing staffing challenges, healthcare systems can achieve better clinical outcomes, improve patient experiences, and support the sustainability of nursing services.

Recommendations for Healthcare Organizations

Based on the findings of this study, the following recommendations are proposed for healthcare organizations:

- Healthcare organizations should regularly assess nurse staffing levels to ensure that staffing matches patient needs and workload requirements.
- Hospitals should develop evidence-based staffing plans that consider patient complexity, nurse qualifications, and clinical demands.
- Healthcare facilities should establish safe nurse-to-patient ratios to reduce excessive workload and improve patient safety.
- Organizations should invest in recruitment and retention programs to address nursing shortages and maintain an adequate workforce.
- Healthcare managers should create supportive working environments that reduce nurse stress, burnout, and job dissatisfaction.
- Hospitals should provide continuous professional development opportunities to improve nurses' knowledge and clinical skills.
- Healthcare organizations should use quality indicators such as patient satisfaction, mortality rates, and adverse events to evaluate the effectiveness of staffing strategies.
- Nursing leadership should participate in staffing decisions to ensure that workforce planning reflects the realities of clinical practice.

for Nursing Practice

The findings of this study provide several recommendations for improving nursing practice

- Nurses should continue to follow evidence-based practices to provide safe and effective patient care despite staffing challenges.
- Nursing teams should maintain effective communication and collaboration to improve patient management and reduce errors.
- Nurses should participate in identifying staffing concerns and reporting situations where workload affects patient safety.
- Healthcare professionals should promote teamwork and appropriate delegation to improve efficiency in clinical settings.
- Experienced nurses should support and mentor less experienced nurses to strengthen the quality of nursing care.
- Nurses should engage in continuous learning and professional development to enhance their clinical competencies.
- Nursing professionals should advocate for safe staffing conditions that support both patient safety and workforce well-being.

Improving nursing practice requires not only individual responsibility but also organizational support. Nurses perform more effectively when they work in environments that provide adequate resources, appropriate staffing, and opportunities for professional growth.

Suggestions for Future Research

Although this study provides important information regarding the impact of nurse staffing levels on patient outcomes and quality of care, further research is recommended to explore this topic in greater depth.

Future studies should examine nurse staffing effects across different healthcare settings, including intensive care units, emergency departments, surgical wards, and community healthcare services. Research comparing staffing models across different countries and healthcare systems may provide additional understanding of effective workforce strategies.

Further research should also investigate the long-term effects of staffing improvements on patient outcomes, nurse retention, and healthcare costs. Future studies may explore

how factors such as nurse education, experience, leadership support, and workplace environment interact with staffing levels to influence quality of care.

Qualitative research involving nurses and patients may provide deeper insight into their experiences regarding staffing challenges and healthcare quality. Additionally, future studies should evaluate innovative staffing approaches, including technology-supported workforce planning and predictive staffing models.

Continued research in this area is necessary to develop effective strategies that address nursing shortages, improve patient safety, and ensure sustainable delivery of high-quality healthcare services.

REFERENCES

Aiken, L. H., Clarke, S. P., Sloane, D. M., Sochalski, J., & Silber, J. H. (2002). Hospital nurse staffing and patient mortality, nurse burnout, and job dissatisfaction. JAMA, 288(16), 1987–1993.

Aiken, L. H., Sloane, D. M., Bruyneel, L., Van den Heede, K., & Sermeus, W. (2014). Nurses' reports of working conditions and hospital quality of care in 12 countries. International Journal of Nursing Studies, 50(2), 143–153.

Blegen, M. A., Goode, C. J., Spetz, J., Vaughn, T., & Park, S. H. (2011). Nurse staffing effects on patient outcomes: Safety-net and non-safety-net hospitals. Medical Care, 49(4), 406–414.

Buchan, J., Duffield, C., & Jordan, A. (2019). Solving nursing shortages: Improving recruitment and retention of nurses. Journal of Advanced Nursing, 75(12), 2923–2934.

Cho, E., Sloane, D. M., Kim, E. Y., Kim, S., Choi, M., Yoo, I. Y., Lee, H. S., & Aiken, L. H. (2017). Effects of nurse staffing, work environments, and education on patient mortality: An analysis of hospital data. International Journal of Nursing Studies, 67, 1–8.

Dall'Ora, C., Ball, J., Reinius, M., & Griffiths, P. (2020). Burnout in nursing: A theoretical review. Human Resources for Health, 18, 41.

Donabedian, A. (1988). The quality of care: How can it be assessed? JAMA, 260(12), 1743–1748.

Griffiths, P., Maruotti, A., Saucedo, A. R., Redfern, O. C., Ball, J. E., Briggs, J., & Dall'Ora, C. (2018). Nurse staffing, nursing assistants and hospital mortality: Retrospective longitudinal cohort study. BMJ Quality & Safety, 28(8), 609–617.

Haegdorens, F., Van Bogaert, P., De Meester, K., & Monsieurs, K. G. (2019). The impact of nurse staffing levels and nursing workload on hospital mortality. Journal of Nursing Management, 27(6), 1233–1244.

Hugonnet, S., Chevrolet, J. C., & Pittet, D. (2007). The effect of workload on infection control practices and healthcare-associated infections. Infection Control & Hospital Epidemiology, 28(3), 264–269.

Irvine, D., Sidani, S., & McGillis Hall, L. (1998). Linking outcomes to nurses' roles in health care. Nursing Economics, 16(2), 58–64.

Kalisch, B. J., Landstrom, G., & Hinshaw, A. S. (2009). Missed nursing care: A concept analysis. Journal of Advanced Nursing, 65(7), 1509–1517.

Kane, R. L., Shamliyan, T. A., Mueller, C., Duval, S., & Wilt, T. J. (2007). The association of registered nurse staffing levels and patient outcomes. Medical Care, 45(12), 1195–1204.

Kutney-Lee, A., McHugh, M. D., Sloane, D. M., Cimiotti, J. P., Flynn, L., Neff, D. F., & Aiken, L. H. (2009). Nursing: A key to patient satisfaction. Health Affairs, 28(4), 669–677.

Lasater, K. B., Aiken, L. H., Sloane, D. M., French, R., Martin, B., Reneau, K., & McHugh, M. D. (2021). Chronic hospital nurse understaffing meets COVID-19: An observational study. BMJ Quality & Safety, 30(8), 639–647.

Lucero, R. J., Lake, E. T., & Aiken, L. H. (2010). Nursing care quality and adverse events in hospitals. Journal of Nursing Administration, 40(6), 274–281.

McHugh, M. D., Kutney-Lee, A., Cimiotti, J. P., Sloane, D. M., & Aiken, L. H. (2011). Nurses' widespread job dissatisfaction, burnout, and frustration with health benefits. Health Affairs, 30(2), 202–210.

Needleman, J., Buerhaus, P., Mattke, S., Stewart, M., & Zelevinsky, K. (2002). Nurse-staffing levels and the incidence of adverse outcomes in hospitalized patients. New England Journal of Medicine, 346(22), 1715–1722.

Needleman, J., Buerhaus, P., Pankratz, V. S., Leibson, C. L., Stevens, S. R., & Harris, M. (2011). Nurse staffing and inpatient hospital mortality. New England Journal of Medicine, 364(11), 1037–1045.

Oldland, E., Botti, M., Hutchinson, A. M., & Redley, B. (2020). A framework of nurses' responsibilities for quality healthcare. Journal of Advanced Nursing, 76(3), 699–707.

Stone, P. W., Mooney-Kane, C., Larson, E. L., Horan, T., Glance, L. G., Zwanziger, J., & Dick, A. W. (2007). Nurse working conditions and patient safety outcomes. Medical Care, 45(6), 571–578.

Stalpers, D., de Brouwer, B. J. M., Kaljouw, M. J., & Schuurmans, M. J. (2015). Associations between characteristics of the nurse work environment and five nurse-sensitive patient outcomes. Applied Nursing Research, 28(4), 285–292.

Tourangeau, A. E., Doran, D. M., McGillis Hall, L., O'Brien Pallas, L., Pringle, D., Tu, J. V., & Cranley, L. A. (2007). Impact of hospital nursing care on 30-day mortality for acute medical patients. Journal of Advanced Nursing, 57(1), 32–44.

World Health Organization. (2020). State of the World's Nursing 2020: Investing in education, jobs and leadership.

Aiken, L. H., Sermeus, W., Van den Heede, K., Sloane, D. M., Busse, R., McKee, M., et al. (2012). Patient safety, satisfaction, and quality of hospital care. BMJ, 344, e1717.

Ball, J. E., Bruyneel, L., Aiken, L. H., Sermeus, W., Sloane, D. M., Rafferty, A. M., et al. (2018). Post-operative mortality, missed care and nurse staffing. International Journal of Nursing Studies, 78, 10–15.

Bae, S. H. (2022). Nurse staffing and patient outcomes: A review of current evidence. Nursing Outlook, 70(2), 227–236.

Cho, S. H., Ketefian, S., Barkauskas, V. H., & Smith, D. G. (2003). The effects of nurse staffing on adverse events, morbidity, mortality, and medical costs. Nursing Research, 52(2), 71–79.

Duffield, C., Diers, D., O'Brien-Pallas, L., Aisbett, C., Roche, M., King, M., & Aisbett, K. (2011). Nursing staffing, nursing workload, the work environment and patient outcomes. Applied Nursing Research, 24(4), 244–255.

Griffiths, P., Ball, J., Drennan, J., Dall'Ora, C., Jones, J., Maruotti, A., & Simon, M. (2016). Nurse staffing and patient outcomes: Strengths and limitations of evidence. Journal of Advanced Nursing, 72(10), 2580–2593.

Needleman, J. (2015). Nurse staffing: The knowns and unknowns. Nursing Economics, 33(1), 5–7.

Rafferty, A. M., Clarke, S. P., Coles, J., Ball, J., James, P., McKee, M., & Aiken, L. H. (2007). Outcomes of variation in hospital nurse staffing in English hospitals. Journal of Nursing Administration, 37(12), 527–535.

Rochefort, C. M., & Clarke, S. P. (2010). Nurses' work environments, workload, and patient outcomes. Nursing Research, 59(1), 45–52.

Shekelle, P. G. (2013). Nurse-patient ratios as a patient safety strategy. Annals of Internal Medicine, 158(5), 404–409.

Shin, S., Park, J. H., & Bae, S. H. (2018). Nurse staffing and hospital outcomes: A systematic review. International Journal of Nursing Studies, 84, 15–22.

IEER
Institute for Excellence in Education

Sloane, D. M., Smith, H. L., McHugh, M. D., & Aiken, L. H. (2018). Effect of changes in nurse staffing on patient outcomes. Health Services Research, 53(4), 2367–2388.

Twigg, D., Gelder, L., & Myers, H. (2015). The impact of nurse staffing methodology on patient outcomes. Journal of Advanced Nursing, 71(4), 801–810.

Van den Heede, K., Clarke, S. P., Sermeus, W., Vleugels, A., & Aiken, L. H. (2007). International experts' perspectives on nurse staffing and patient outcomes. Journal of Advanced Nursing, 60(3), 289–297.

Zúñiga, F., Ausserhofer, D., Hamers, J. P. H., Engberg, S., & Schwendimann, R. (2015). Are staffing, work environment, and missed care associated with patient outcomes? International Journal of Nursing Studies, 52(10), 1634–1643.

IEER
Institute for Excellence in Education & Research