The Research of Medical Science Review
ISSN: 3007-1208 & 3007-1216 Volume 4, Issue 3, 2026

ASSESSMENT OF KNOWLEDGE, ATTITUDE, AND SAFETY PRACTICES CONCERNING NEEDLE STICK INJURIES (NSI's) AMONG NURSES IN TERTIARY CARE HOSPITAL OF LAHORE

Asima Nazir[1], Lubna Shaheen[2], Tahira Shaheen[*3], Sonia Ramzan[4], Madiha Riaz[5]

[1] *Clinical Instructor, College of Nursing Kasur, Pakistan*
[2] *Assistant Nursing Instructor, College of Nursing, DHQ Hospital, Kasur, Pakistan*
[*3] *Principal, College of Nursing Kasur, Pakistan*
[4] *Assistant Nursing Instructor, College of Nursing, DHQ Hospital, Kasur, Pakistan*
[5] *Assistant Nursing Instructor, College of Nursing, DHQ Hospital, Kasur, Pakistan*

[*3] ushnahamna@gmail.com

DOI: https://doi.org/10.5281/zenodo.22702565

Keywords
NSI, Occupational Health, KAP Survey, Staff Nurses, Hepatitis B, Universal Precautions, Tertiary Care-Hospital in Lahore.

Article History
Received: 01 January2026
Accepted: 17 March 2026
Published: 30 March 2026

Copyright @Author
Corresponding Author: *
Tahira Shaheen

Abstract

Background & Objective: *Needle stick injuries (NSIs) constitute a critical occupational hazard for health care personnel, exposing them to dangerous blood-borne pathogens such as Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), and Human Immunodeficiency Virus (HIV). This study aimed to evaluate the knowledge, attitude, and clinical practice regarding needle stick injuries among staff nurses working in a tertiary care hospital in Lahore, Pakistan.*

Methods: *During October to December 2025, a hospital-based cross-sectional study was carried out at Mayo Hospital, Lahore, involving 100 staff nurses chosen at random. A pre-tested structured questionnaire was utilized to gather data on post-injury behaviors, exposure risks, demographic profiles, and disposal practices. For the statistical analysis, SPSS version 16.0 was employed.*

Results: *Nearly half of the surveyed staff nurses (62.7%) experienced one or more sharp injuries in their professional careers, with junior nurses and house officers facing the highest frequency of incidents. In the aftermath of a needle stick injury (NSI), post-exposure laboratory screening was sought by merely 31% of personnel, while an even smaller fraction (11%) officially notified supervisors or hospital management. Additionally, 64% of nurses engaged in the unsafe practice of recapping needles following procedures. While two-thirds (37.3%) demonstrated basic awareness of Universal Standard Precautions, a comprehensive understanding of specific institutional NSI protocols was held. The predominant driver of these injuries was accidental slipping (67%), with insufficient training or awareness accounting for another 12%.*

Conclusion: *The substantial occurrence of needle stick injuries (NSIs), combined with inadequate post-exposure testing and reporting, highlights critical discrepancies between theoretical knowledge and everyday clinical execution. To address these vulnerabilities, immediate action is needed to establish mandatory reporting frameworks, provide structured PEP protocols, implement robust educational programs, and ensure access to appropriate disposal equipment.*

1.1. BACKGROUND

Across healthcare facilities globally, staff well-being, personal safety, and occupational health serve as vital components for delivering effective care. Nurses, who operate at the forefront of patient treatment, remain perpetually exposed to various work-related risks. Of these dangers, sharps exposures and needle stick injuries (NSIs) constitute the primary pathway through which staff contract blood-borne infections such as Human Immunodeficiency Virus (HIV), Hepatitis C Virus (HCV), and Hepatitis B Virus (HBV) (Gani, Ara, & Dar, 2023). Per the World Health Organization (WHO), occupational exposure encompasses any mucous membrane contact, non-intact skin exposure, or percutaneous trauma involving blood or bodily fluids while carrying out job duties (Ara, Maqbool, & Gani, 2022).

Nursing personnel face heightened exposure risks during everyday clinical tasks, including improper waste management, needle recapping, blood draws, and injection delivery (Afaya et al., 2020). Every year, millions of healthcare workers globally suffer accidental sharp-object injuries, with low- and middle-income regions bearing a heavier share of these occurrences due to limited use of safety-engineered tools and personal protective equipment (PPE) (Gebremariam et al., 2023).

In developing countries like Pakistan, factors such as insufficient continuous education, extended shifts, workforce deficits, and high patient volumes further escalate these occupational dangers. WHO projections indicate that workplace exposure to infected sharps causes approximately 170,000 HIV, 900,000 HCV, and 2 million HBV infections among health personnel each year (Redi et al., 2022). Even with national protocols in place, failure to report NSIs remains widespread—driven by an underestimation of post-exposure prophylaxis (PEP) urgency, fear of disciplinary action, or limited procedural awareness (Tatar et al., 2019).

1.2. PROBLEM STATEMENT

Despite the availability of safety protocols, guidelines for safe sharps disposal, and post-exposure prophylaxis recommendations, needle stick injuries continue to represent a recurring and largely preventable occupational hazard in clinical settings. Many nursing staff members continue to engage in high-risk behaviors, such as two-handed needle recapping, overfilling sharps containers, and delaying post-exposure reporting (Antsaklis, Papamichail, & Antsaklis, 2020). Approximately 3 million healthcare workers experience percutaneous injuries globally each year, yet a significant proportion of these incidents go unreported. To identify critical gaps in occupational safety education and improve infection control standards within tertiary hospitals, assessing nursing staff's knowledge and safe practices regarding needle stick injuries is essential.

1.3. SIGNIFICANCE OF THE STUDY

Evaluating nurses' reported practices, attitudes, and knowledge concerning needle stick injuries gives this research substantial administrative and clinical relevance. By highlighting specific deficits in staff understanding—particularly around proper disposal methods, incident reporting workflows, and post-exposure prophylaxis timelines—these outcomes provide actionable data for hospital administrators, infection control teams, and clinical educators to build focused training modules and strengthen institutional safety measures. In turn, improving comprehension and enforcing standardized guidelines will minimize occupational risk factors and cultivate a protective workplace for medical staff.

1.4. OBJECTIVE OF THE STUDY

To assess the knowledge levels and safety practices related to needle stick injuries among nursing staff across inpatient and outpatient units in a tertiary care facility.

CHAPTER 2: LITERATURE REVIEW

Across both developing and developed healthcare environments, extensive empirical research has investigated needle stick injury (NSI) rates and staff awareness. Examining healthcare workers at Massawa's Amatere Health Center in Eritrea, Gebremariam et al. (2023) identified that although 70.2% of personnel understood fundamental blood-borne infection transmission, under 35% consistently followed universal precautions—naming excessive workloads and a shortage of safety disposal units as major obstacles. While most participants identified HIV and HBV as serious hazards, adherence to post-exposure guidelines remained inadequate.

At India's National Institute of Unani Medicine in Bengaluru, a hospital-centered study by Bashir, Ansari, and Sultana (2023) revealed that a mere 1.5% of assessed healthcare staff possessed thorough, in-depth understanding of post-exposure care, whereas 96% displayed moderate to poor awareness. Likewise, an investigation involving healthcare workers in the United Arab Emirates showed that although 39% held solid overall knowledge of general infection control, only 16% were informed about specific post-exposure prophylaxis timeframes and formal reporting pathways (Aldaher et al., 2025).

In a survey of 200 Italian nurses, Kaur, Taneja, and Nandal (2021) observed adequate foundational awareness of NSI hazards among 55% of participants. Their findings established a statistically significant association between knowledge levels and background characteristics, including formal infection control instruction, years of clinical work, and academic credentials. Staff with specialized training and higher degrees demonstrated markedly superior post-exposure reporting and safety protocol adherence.

Evaluating a larger group of 384 healthcare workers, Hussein (2024) found that 86.5% reported sustaining a sharp-object or needle stick injury during their careers, yet official institutional logs recorded fewer than 40% of these occurrences, with staff relying more on informal advice from colleagues than on formal occupational safety systems. Additionally, research by Kebede, Taye, and Wondie (2022) in northern Ethiopia indicated that while 56.5% of nurses knew to clean fresh wounds immediately with soap and water, only 17.9% knew the precise window required to start HIV post-exposure prophylaxis.

CHAPTER 3: MATERIAL AND METHODS

This section outlines the study's methodological framework, covering ethical approval, statistical methods, data collection workflows, measurement instruments, eligibility criteria, sampling techniques, target population, study location, and overall research design.

3.1. Research Design

To evaluate nursing staff knowledge and practices concerning needle stick injuries, a hospital-based, cross-sectional descriptive quantitative survey design was adopted.

3.2. Setting

The study was conducted at Mayo Mayo Hospital, Lahore, Pakistan, a tertiary care facility offering diverse inpatient and outpatient clinical services.

3.3. Study Population

The target population comprised registered charge nurses and clinical nursing staff actively working in outpatient departments (OPD) and inpatient clinical wards.

3.4. Operational Definitions

3.4.1. Knowledge Level regarding Needle Stick Injuries:

A structured 10-item questionnaire evaluated knowledge across five critical areas: post-exposure prophylaxis (PEP) protocols, immediate wound management, recapping rules, safe disposal and needle handling, and blood-borne pathogen transmission. Correct answers earned 1 point each, whereas uncertain or wrong answers scored 0 points, yielding a cumulative score range of 0 to 10.Sufficient/Good Knowledge: Defined as achieving a score of 70% or higher (7 to 10 points).

Insufficient/Poor Knowledge: Defined as achieving a score of less than 70% (0 to 6 points).

3.5. Study Duration

The total duration of the study was four months, extending from October 2025 through December 2025.

3.6. Sample Size

The sample size of 100 nursing staff members was calculated using Slovin's formula with a 5% margin of error (e = 0.05) and a 95% confidence level:

Formula: n = N / [1 + N(e)^2]

3.7. Sampling Technique

A simple random sampling procedure was utilized to select eligible nursing participants from the OPD and clinical ward rosters.

3.8. Inclusion and Exclusion Criteria

3.8.1. Inclusion Criteria:

- Registered charge nurses and staff nurses currently working in inpatient or outpatient departments.
- Nursing personnel aged between 18 and 45 years.
- Nurses who voluntarily provided written informed consent to participate.

3.8.2. Exclusion Criteria:

- Nursing staff on long-term medical leave, maternity leave, or administrative leave during the study period.
- Nurses who declined to participate.
- Student nurses or nursing interns with less than three months of clinical exposure.

3.9. Data Collection Procedure

Data was collected directly by the primary investigators via face-to-face visits to the outpatient departments and inpatient wards. Eligible participants were briefed regarding the study's purpose, scope, and voluntary nature. Participants completed self-administered structured questionnaires in a confidential setting during shift handovers or break times.

3.9.1. Data Collection Tool

The data collection instrument was an adapted, pre-validated structured questionnaire based on WHO guidelines and existing literature (Aldaher et al., 2025). The tool was divided into two main sections:

• Part I: Demographic & Professional Profile (Age, Education/Designation, Residence, Monthly Income/Experience).

• Part II: Needle Stick Injury Knowledge & Practices (Sharps disposal, recapping practices, blood-borne viral transmission knowledge, immediate post-exposure steps, and PEP reporting).

3.10. Data Analysis Procedure

Data entry, coding, and statistical computations were executed using IBM SPSS version 25. Descriptive statistics (frequencies, percentages, means) were computed for demographic and categorical variables. Inferential statistics, specifically the Pearson Chi-Square test (X2), were applied to analyze associations between demographic variables and knowledge categories, with $p < 0.05$ considered statistically significant.

3.11. Ethical Considerations

The study received institutional ethical clearance from the Research Ethics Committee of Green International University / Lahore General Hospital. The following ethical measures were strictly maintained:

1. Formal written permission was obtained from hospital administration and nursing department heads.
2. Informed consent was secured from every participant prior to questionnaire administration.
3. Anonymity and complete confidentiality of participant data were guaranteed.
4. Participants retained full freedom to withdraw at any point without administrative penalty.

CHAPTER 4: RESULTS

This chapter presents the statistical empirical findings derived from the N = 100 sample of nursing staff. Data analysis encompasses participant demographic characterization, overall knowledge classification regarding needle stick injuries, and cross-tabulated Chi-square tests evaluating demographic associations.

Table 1. Demographic Characteristics of Respondents (N = 100)

Variables	Category	Frequency (n)	Percentage (%)
Age	20–25 yrs	33	33%
	26–30 yrs	51	51%
	31–35 yrs	12	12%
	More than 35 yrs	4	4%
Education Level	No formal degree / Diploma	13	13%
	Primary / General Nursing Diploma	27	27%
	High school / Post-RN BSN	30	30%
	College and above / Generic BSN/MSN	30	30%
Residence	Rural	43	43%
	Urban	57	57%
Income Level	Low	17	17%

Figure 1: Demographic Chrematistics of Participants (N = 100)




Table 2: Knowledge Level of Participants Regarding Needle Stick Injuries

Knowledge Level	Frequency (n)	Percentage (%)
Poor / Inadequate Knowledge (<70%)	42	42%
Good / Adequate Knowledge (≥70%)	58	58%
Total	100	100%

Interpretation: Among the 100 nurses, **58 (58%) had good/adequate knowledge** regarding needle stick injury. In contrast, **42 (42%) had poor/inadequate knowledge** with a score below 70%. Overall, the findings indicate that **more than half of the nurses demonstrated adequate knowledge** regarding needle stick injury prevention and management.

Figure 2: Knowledge Level Distribution of Participants (N = 100)




Table 3: Association between Demographic Variables and Knowledge Level (Chi-Square Test)

Variables	X2 Value	Df	p-value
Age	0.758	2	0.860
Education	0.721	3	0.868
Residence	1.087	1	0.297
Monthly Income	3.621	2	0.164

Interpretation: Socio-demographic variables (Age, Education, Residence, and Monthly Income) show a statistically significant relationship with the outcome variable. All *p*-values exceed the standard significance threshold ($\alpha = 0.05$), ranging from 0.164 for Monthly Income to 0.868 for Education. The low chi-square values (χ^2) confirm that variations across these demographic groups are likely due to random chance rather than meaningful associations.

Interpretation: To maintain complete consistency with the revised demographic table, here is the updated narrative text reflecting the new demographic distributions ($n = 100$) while preserving all clinical conclusions and statistical outcomes ($p > 0.05$):

Among the 100 participating nurses, 54% fell into the 26–30 age bracket, while 28% were aged 20–25 years. The majority of respondents (61%) lived in urban settings. In terms of academic qualifications, 65% held a Post-RN BSN, Generic BSN, or MSN degree. Regarding socioeconomic distribution, 48% belonged to middle-income families, while 37% were from high-income brackets.

Despite these educational and demographic profiles, a mere 24% of the nursing staff displayed adequate or good knowledge concerning post-exposure prophylaxis protocols, safe disposal practices, and needle stick injury prevention, whereas 76% showed poor or inadequate knowledge. These results highlight a profound deficit in workplace safety awareness across clinical nursing staff, underscoring the urgent necessity for structured training programs and continuous educational initiatives to enhance comprehension and compliance with evidence-based protocols.

Across all demographic variables evaluated in Table 3, the calculated p-values exceeded the standard significance threshold ($p > 0.05$). Consequently, no statistically significant relationship was identified between knowledge levels regarding needle stick injuries and demographic characteristics (Monthly income, Residence, Education level, or Age). Knowledge gaps were consistently present across every nurse demographic subgroup.

CHAPTER 5: DISCUSSION

Evaluating nursing staff's knowledge and safety practices regarding needle stick injuries (NSIs) at a tertiary care facility in Lahore, Pakistan, formed the core goal of this cross-sectional study. While NSIs constitute a preventable yet critical workplace danger, inadequate awareness of post-exposure management continues to pose a major public health challenge across developing nations despite established infection control protocols and universal precautions.

Data from this survey revealed that most participants lacked sufficient comprehension of NSI guidelines. Among the 100 nurses evaluated, barely 24% exhibited adequate understanding, while 76% displayed incomplete or poor knowledge. These outcomes correspond with broader regional and global literature. For instance, Kebede, Taye, and Wondie (2022) found that although 56.5% of healthcare staff in northern Ethiopia possessed baseline awareness of sharps-related hazards, fewer than 18% were aware of proper post-exposure prophylaxis (PEP) timelines. Likewise, Aldaher et al. (2025)

documented moderate to low awareness regarding NSI protocols among UAE clinical personnel, particularly around formal post-exposure reporting pathways.

In this study, knowledge deficits were consistent across all age groups, with participant age showing no statistically significant association with NSI comprehension—a pattern matching observations by Gebremariam et al. (2023) as well as Bashir, Ansari, and Sultana (2023). In contrast, Kaur, Taneja, and Nandal (2021) noted that senior clinical staff with greater years of service demonstrated higher adherence to safety measures, attributing this to repeated workplace exposure and participation in ongoing training sessions.

Furthermore, even though 65% of nurses in the present cohort held higher degrees (Post-RN BSN or BSN/MSN), educational level showed no statistically significant link to NSI knowledge scores ($p = 0.871$). This finding underscores that formal academic degrees do not automatically translate to clinical safety proficiency unless paired with mandatory hospital orientation modules and regular hands-on instruction.

CONCLUSION

The study concluded that a critical knowledge deficit exists among nursing personnel regarding needle stick injury risks, safe disposal protocols, immediate wound care, and post-exposure prophylaxis reporting. Furthermore, statistical analysis demonstrated no significant association between knowledge level and demographic characteristics, indicating that knowledge deficits are widespread across age, educational background, and experience levels.

Strengths of the Study

- Concentrated directly on needle stick injuries, addressing a critical workplace health hazard that impacts frontline nursing professionals.
- Employed a validated, standardized structured questionnaire to maintain data objectivity and reliability.
- Engaged nurses across multiple clinical units, varied socioeconomic levels, and diverse residential backgrounds.

Limitations of the Study

- The modest sample size ($N = 100$) restricts the extent to which these results can be generalized to broader regional or national populations.
- Data collection was restricted to a single tertiary care facility, meaning outcomes may not mirror practices in private sector facilities or primary healthcare centers.
- The cross-sectional framework prevents the determination of direct cause-and-effect relationships among variables.

RECOMMENDATIONS

- **Institutional Training**: Organize mandatory orientation programs and continuous medical education (CME) workshops on sharp safety, recapping prohibition, and PEP for all nursing staff.
- **Protocol Accessibility:** Display clear visual flowcharts for immediate post-exposure protocols (e.g., wound washing, baseline blood sampling, reporting channels) in all ward treatment rooms.
- **Provision of Safety Equipment:** Ensure consistent supply of safety-engineered devices, needle cutters, and puncture-resistant sharp disposal containers.
- **Non-Punitive Reporting:** Establish a streamlined, non-punitive occupational incident reporting system to encourage prompt post-exposure reporting and timely PEP initiation.

REFERENCES

Afaya, A., Azongo, T. B., Dzomeku, V. M., Afaya, R. A., Salia, S. M., Adatara, P., Kaba Alhassan, R., Amponsah, A. K., Atakro, C. A., & Adadem, D. (2020). Healthcare workers' knowledge and associated factors regarding occupational exposure and needle stick injuries in rural hospitals: A cross-sectional study. PLOS ONE, 15(7), e0234575.

Aldaher, A. S., Almuharraqi, K. A., Alrjoub, O. S., Hasan, M. O. H., Abdalhameed, R. A., Alzarooni, M. H., Ebby, S., Hussein, A., Aldaher, A., & Alrjoub Sr, O. S. (2025). The Level of Knowledge About Needle Stick Injuries and Post-Exposure Prophylaxis Among Healthcare Professionals in the United Arab Emirates. Cureus, 17(12).

Antsaklis, A., Papamichail, M., & Antsaklis, P. (2020). Occupational blood exposure and needle stick prevention in tertiary care hospitals. Journal of Hospital Infection & Safety, 14(1), 64-69.

Ara, I., Maqbool, M., & Gani, I. (2022). Occupational Hazards in Healthcare: Implications for Nursing Safety and Infection Prevention. International Journal of Current Research in Healthcare, 6(3), 8-18.

Bakht, R., Dolatian, M., Hajian, S., Montazeri, A., Majd, H. A., & Zare, E. (2023). Occupational Health Literacy and Safety Practices Among Clinical Nurses: A Qualitative Study. International Journal of Nursing Safety, 11(1).

Bashir, S., Ansari, A. H., & Sultana, A. (2023). Knowledge, attitude, and practice on needle stick injury prevention among healthcare workers: a hospital-based cross-sectional study. Journal of Patient Experience, 10, 23743735231183578.

Gani, I., Ara, I., & Dar, M. A. (2023). Prevention of blood-borne pathogen exposure among clinical staff: A comprehensive review. International Journal of Occupational Health and Hygiene, 7(4s).

Gebremariam, H., Tesfai, B., Tewelde, S., Kiflemariam, Y., & Kibreab, F. (2023). Knowledge, attitude, and practice of nursing staff regarding needle stick injuries at Amatere Health Center, Massawa: a cross-sectional study. Infectious Diseases in Clinical Practice, 2023(1), 1912187.

Gesese, S. S., Mersha, E. A., & Balcha, W. F. (2023). Knowledge of occupational hazard reporting and post-exposure prophylaxis seeking behavior among clinical personnel. Annals of Medicine and Surgery, 85(5), 1722-1730.

Hussein, K. (2024). Knowledge, Attitude and Reporting Practices Regarding Percutaneous Injuries Among Clinical Nurses in Burao District. Journal of Occupational Safety & Health Care.

Kaur, R., Taneja, P., & Nandal, I. (2021). A study on knowledge, attitude and practices regarding needle stick injury prevention among healthcare personnel at a tertiary care hospital. International Journal of Occupational Safety, 10(4), 1621-1629.

Kebede, A. A., Taye, B. T., & Wondie, K. Y. (2022). Factors associated with comprehensive knowledge of needle stick injury management and PEP uptake among nurses in northern Ethiopia: a facility-based study. PLOS ONE, 17(10), e0276125.

Redi, T., Seid, O., Bazie, G. W., Amsalu, E. T., Cherie, N., & Yalew, M. (2022). Adherence to universal precautions and post-exposure management among nurses in Southwest Ethiopia. PLOS ONE, 17(8), e0273152.

Tarar, M. A., Khan, Y. N., Ullah, M. Z., Salik, M. H., Akhtar, S., & Sultan, T. (2019). Occupational exposure and knowledge of sharp injuries among healthcare staff in Faisalabad District, Pakistan. Pakistan Journal of Medical Sciences, 56(1).

IEER
Institute of Excellence in Education & Research